

Name:

Period:

Seat#:

Directions: Show all work in a way that would earn you credit on the AP Test! This is always the rule! Answers provided at the bottom of the last page. **SHOW WORK ON BINDER PAPER. STAPLE BINDER PAPER TO THE BACK OF THIS WORKSHEET!** Clearly label which question number your work is for.

NChO 1999

Reaction	ΔH
$Mg(s) + 2 HCl(aq) \rightarrow MgCl_2(aq) + H_2(g)$	-467 kJ mol ⁻¹
$MgO(s) + 2 HCl(aq) \rightarrow MgCl_2(aq) + H_2O(l)$	-151 kJ mol ⁻¹

27. According to this information, and given the fact that for water, $\Delta H_f = -286$ kJ mol⁻¹, what is ΔH_f for MgO(s)?
- (A) -904 kJ mol⁻¹ (C) -334 k J mol⁻¹
 (B) -602 kJ mol⁻¹ (D) -30 kJ mol⁻¹

NChO 1998


22. Carbon reacts with oxygen according to this equation.
 $2C(s) + O_2(g) \rightarrow 2CO(g) \quad \Delta H = -220$ kJ

Which statements are true?

1. The reaction is exothermic.
 2. The combustion of 0.50 mol of carbon produces 55 kJ

Substance	ΔH°_f (kJ mol ⁻¹)	of heat energy
Fe ₂ O ₃ (s)	-824.2	(A) 1 only
Fe(OH) ₃ (s)	-823.0	(C) both 1 and 2
H ₂ O(l)	-285.8	(B) 2 only (D) neither 1 nor 2

24. Use these data to calculate ΔH° for this reaction.



Reaction	ΔH° , kJ mol ⁻¹
$\frac{1}{2} N_2(g) + \frac{1}{2} O_2(g) \rightarrow NO(g)$	90.2 kJ mol ⁻¹
$\frac{1}{2} N_2(g) + O_2(g) \rightarrow NO_2(g)$	33.2 kJ mol ⁻¹

25. A 1.0 g sample of substance A at 100 °C is added to 100 mL of H₂O at 25 °C. Using separate 100 mL portions of H₂O, the procedure is repeated with substance B and then with substance C. How will the final temperatures of the water compare?

- (A) T_c > T_b > T_a (C) T_a > T_b > T_c
 (B) T_b > T_a > T_c (D) T_a = T_b = T_c


Substance	Specific Heat
A	0.60 J g ⁻¹ °C ⁻¹
B	0.40 J g ⁻¹ °C ⁻¹
C	0.20 J g ⁻¹ °C ⁻¹

26. How many grams of benzene, C₆H₆(l), must be burned in a bomb calorimeter to raise its temperature by 1.5 °C? Given: The calorimeter constant is 12.59 kJ C⁻¹ and the $\Delta H^\circ_{\text{combustion}}$ for C₆H₆ = -41.9 kJ g⁻¹
- (A) 0.45 g (C) 3.3 g
 (B) 2.8 g (D) 8.4 g

NChO 1997

19. 30.0 mL of water at 10. °C is mixed with 50.0 mL of water at 60. °C. What is the final temp of the mixture?
- (A) 31 °C (C) 41 °C
 (B) 35 °C (D) 46 °C
24. What is the value of ΔH° for this reaction?
- Fe₂O₃(s) + 3H₂O(l) → 2Fe(OH)₃(s)
- (A) 35.6 kJ (C) 858.6 kJ
 (B) 286 kJ (D) -536 kJ

25. When Na₂S₂O₃ · 3H₂O dissolves in water, the solution gets cold. Which energy diagram best represents the behavior of this solution process?



NChO 1996

22. The standard enthalpy of formation (ΔH°_f) for sodium bromide is the enthalpy change for the reaction
- (A) Na⁺(g) + Br⁻(g) → NaBr(g)
 (B) Na⁺(g) + Br⁻(g) → NaBr(s)
 (C) 2 Na(s) + Br₂(g) → 2 NaBr(s)
 (D) Na(s) + ½ Br₂(l) → NaBr(s)
23. Use the standard enthalpies of formation to calculate ΔH° for this reaction:
- 2 CrO₄²⁻(aq) + 2 H⁺(aq) → Cr₂O₇²⁻(aq) + H₂O(l)
- (A) 272.1 kJ (C) -13.7 kJ
 (B) 13.7 kJ (D) -272.1 kJ


Substance	ΔH°_f , kJ mol ⁻¹
CrO ₄ ²⁻ (aq)	- 881.2
Cr ₂ O ₇ ²⁻ (aq)	- 1490.3
H ⁺ (aq)	0
H ₂ O(l)	- 285.8

NChO 1995


21. For which of these processes is the sign of the enthalpy change different from the others?
- (A) $\text{Al}_2\text{O}_3(\text{s}) \rightarrow 2 \text{ Al}(\text{s}) + 3/2 \text{ O}_2(\text{g})$
 (B) $\text{H}_2\text{O}(\text{s}) \rightarrow \text{H}_2\text{O}(\text{l})$
 (C) $\text{Cl}_2(\text{g}) \rightarrow 2\text{Cl}(\text{g})$
 (D) $\text{Cl}(\text{g}) + \text{e}^- \rightarrow \text{Cl}^-(\text{g})$
22. The standard enthalpy of formation (ΔH°_f) for nitrogen(IV) oxide is the enthalpy change for the reaction
 (A) $\text{N}(\text{g}) + 2\text{O}(\text{g}) \rightarrow \text{NO}_2(\text{g})$
 (B) $\frac{1}{2}\text{N}_2(\text{g}) + \text{O}_2(\text{g}) \rightarrow \text{NO}_2(\text{g})$
 (C) $\frac{1}{2}\text{N}_2\text{O}_4(\text{g}) \rightarrow \text{NO}_2(\text{g})$
 (D) $\text{NO}(\text{g}) + \frac{1}{2}\text{O}_2 \rightarrow \text{NO}_2(\text{g})$
23. In a bomb calorimeter, reactions are carried out at
 (A) constant volume. (C) 1 atm pressure and 25 °C.
 (B) constant pressure. (D) 1 atm pressure and 0 °C.
26. Consider the reaction
 $\text{Hg}(\text{l}) + 2 \text{Ag}^+(\text{aq}) \rightarrow \text{Hg}^{2+}(\text{aq}) + 2 \text{Ag}(\text{s})$
 What is enthalpy change for this rxn if ΔH°_f for $\text{Ag}^+(\text{aq})$ is +105.6 kJ mol⁻¹ and for $\text{Hg}^{2+}(\text{aq})$ is +171.1 kJ mol⁻¹?
 (A) 65.5 kJ are evolved per mole of Hg.
 (B) 65.5 kJ are absorbed per mole of Hg.
 (C) 40.1 kJ are evolved per mole of Hg.
 (D) 40.1 kJ are absorbed per mole of Hg.

NChO 1994

24. A student mixes 100 mL of 0.50 M NaOH with 100 mL of 0.50 M HCl in a styrofoam cup and observes a temperature increase of ΔT_1 . When she repeats this experiment using 200mL of each solution, she observes a temperature change of ΔT_2 . If no heat is lost to the surroundings or absorbed by the styrofoam cup, what is true about ΔT_1 and ΔT_2 ?
- (A) $\Delta T_2 = \Delta T_1$ (C) $\Delta T_2 = 2 \Delta T_1$
 (B) $\Delta T_2 = 0.5 \Delta T_1$ (D) $\Delta T_2 = 4 \Delta T_1$
27. Given these values of ΔH° : ΔH°



What is the value of ΔH° for this reaction?

**NChO 1993**

13. Which process or reaction has a positive ΔH ?
- (A) $\text{H}_2\text{O}(\text{l}) \rightarrow \text{H}_2\text{O}(\text{s})$
 (B) $2\text{CH}_3\text{OH}(\text{l}) + 3\text{O}_2(\text{g}) \rightarrow 2\text{CO}_2(\text{g}) + 4\text{H}_2\text{O}(\text{l})$
 (C) $\text{CO}_2(\text{s}) \rightarrow \text{CO}_2(\text{g})$
 (D) $2 \text{Na}(\text{s}) + \text{Cl}_2(\text{g}) \rightarrow 2 \text{NaCl}(\text{s})$
15. For the reaction $\text{H}_2(\text{g}) + \text{I}_2(\text{s}) \rightarrow 2 \text{HI}(\text{g})$ $\Delta H_{\text{rxn}} = 53.0 \text{ kJ}$
 What will be the value of ΔH_{rxn} (in kJ) for this rxn ?
 $\text{HI}(\text{g}) \rightarrow \frac{1}{2} \text{H}_2(\text{g}) + \frac{1}{2} \text{I}_2(\text{s})$
 (A) 26.5 (C) -26.5
 (B) 7.3 (D) -53.0

Answers

1999
 27) B

1998
 22) C
 24) A
 25) C
 26) A

1997
 19) C
 24) A
 25) A

1996
 22) D
 23) C

1995
 21) D
 22) B
 23) A
 26) C

1994
 24) A
 25) C

1993
 13) C
 15) C