# <u>N37 – Acid Base</u>

## Weak Problems

# **N37 – Acid Base** Weak Problems

Target: I can review Acid Base definitions and perform pH calculations for weak acids/bases

# <u>N37 – Acid Base</u>

## Yay, ICE Tables are back!

## **A Weak Acid Equilibrium Problem**

What is the pH of a 0.50 M solution of acetic acid,  $HC_2H_3O_2$ ,  $K_a = 1.8 \times 10^{-5}$ ?

**Step #1:** Write the dissociation equation

 $HC_2H_3O_2 \leftrightarrows C_2H_3O_2^- + H^+$ 

## <u>A Weak Acid Equilibrium Problem</u>

What is the pH of a 0.50 M solution of acetic acid,  $HC_2H_3O_2$ ,  $K_a = 1.8 \times 10^{-5}$ ?

#### Step #2: ICE it!

 $HC_{2}H_{3}O_{2} \leftrightarrows C_{2}H_{3}O_{2}^{-} + H^{+}$   $I \quad 0.50 \quad 0 \quad 0$   $C \quad - \times \quad + \times \quad + \times$   $E \quad 0.50 - \times \quad \times \quad \times \quad \times$ 

## <u>A Weak Acid Equilibrium Problem</u>

What is the pH of a 0.50 M solution of acetic acid,  $HC_2H_3O_2$ ,  $K_a = 1.8 \times 10^{-5}$ ?

**Step #3:** Set up the law of mass action

 $HC_{2}H_{3}O_{2} \leftrightarrows C_{2}H_{3}O_{2}^{-} + H^{+}$   $0.50 - x \qquad x \qquad x$   $1.8 x 10^{-5} = \frac{(x)(x)}{(0.50 - x)} \cong \frac{x^{2}}{(0.50)}$ 

Can use the 5% rule because K < 1 and K at least 1000 time smaller than [initial]

## **A Weak Acid Equilibrium Problem**

What is the pH of a 0.50 M solution of acetic acid,  $HC_2H_3O_2$ ,  $K_a = 1.8 \times 10^{-5}$ ?

**Step #4:** Solve for x, which is also [H<sup>+</sup>]

 $HC_{2}H_{3}O_{2} \leftrightarrows C_{2}H_{3}O_{2}^{-} + H^{+}$ E 0.50 - x x x

$$1.8 x 10^{-5} = \frac{x^2}{(0.50)} \quad [H^+] = 3.0 x 10^{-3} M$$

## **A Weak Acid Equilibrium Problem**

What is the pH of a 0.50 M solution of acetic acid,  $HC_2H_3O_2$ ,  $K_a = 1.8 \times 10^{-5}$ ?

Step #5: Convert [H+] to pH

 $pH = -log(3.0 \ x \ 10^{-3}) = 2.52$ 

### **Reaction of Weak Bases with Water**

The base reacts with water, producing its conjugate acid and hydroxide ion:

 $CH_3NH_2 + H_2O \leftrightarrows CH_3NH_3^+ + OH^- K_b = 4.38 \times 10^{-4}$ 

$$K_{b} = 4.38 x 10^{-4} = \frac{[CH_{3}NH_{3}^{+}][OH^{-}]}{[CH_{3}NH_{2}]}$$

#### K<sub>b</sub> for Some Common Weak Bases

Many students struggle with identifying weak bases and their conjugate acids. What patterns do you see that may help you?

Base	Formula	Conjugate Acid	K <sub>b</sub>
Ammonia	NH <sub>3</sub>	NH <sub>4</sub> +	1.8 x 10 <sup>-5</sup>
Methylamine	CH <sub>3</sub> NH <sub>2</sub>	CH <sub>3</sub> NH <sub>3</sub> +	4.38 x 10 <sup>-4</sup>
Ethylamine	$C_2H_5NH_2$	C <sub>2</sub> H <sub>5</sub> NH <sub>3</sub> <sup>+</sup>	5.6 x 10 <sup>-4</sup>
Diethylamine	$(C_2H_5)_2NH$	$(C_2H_5)_2NH_2^+$	1.3 x 10 <sup>-3</sup>
Triethylamine	(C <sub>2</sub> H <sub>5</sub> ) <sub>3</sub> N	$(C_2H_5)_3NH^+$	4.0 x 10 <sup>-4</sup>
Hydroxylamine	HONH <sub>2</sub>	HONH <sub>3</sub> +	1.1 x 10 <sup>-8</sup>
Hydrazine	$H_2NNH_2$	H <sub>2</sub> NNH <sub>3</sub> +	3.0 x 10 <sup>-6</sup>
Aniline	$C_6H_5NH_2$	C <sub>6</sub> H <sub>5</sub> NH <sub>3</sub> +	3.8 x 10 <sup>-10</sup>
Pyridine	C <sub>5</sub> H <sub>5</sub> N	C₅H₅NH⁺	1.7 x 10 <sup>-9</sup>

## **Reaction of Weak Bases with Water**

The generic reaction for a base reacting with water, producing its conjugate acid and hydroxide ion:

 $\mathbf{B} + \mathbf{H}_{2}\mathbf{O} \leftrightarrows \mathbf{B}\mathbf{H}^{+} + \mathbf{O}\mathbf{H}^{-}$  $K_{b} = \frac{[BH^{+}][OH^{-}]}{[B]}$ 

Yes, all weak bases do this – DO NOT make this more complicated then it needs to be.

What is the pH of a 0.50 M solution of ammonia,  $NH_3$ ,  $K_b = 1.8 \times 10^{-5}$ ?

**Step #1:** Write the equation for the reaction

 $NH_3 + H_2O \leftrightarrows NH_4^+ + OH^-$ 

What is the pH of a 0.50 M solution of ammonia,  $NH_3$ ,  $K_b = 1.8 \times 10^{-5}$ ?

Step #2: ICE it!

	NH <sub>3</sub> +	$H_2O \leftrightarrows NH_4^+$	+ OH <sup>-</sup>
Ι	0.50	0	0
С	- X	+X	+X
E	0.50 - x	×	×

What is the pH of a 0.50 M solution of ammonia,  $NH_3$ ,  $K_b = 1.8 \times 10^{-5}$ ?

**Step #3: Set up the law of mass action** 

 $NH_{3} + H_{2}O \leftrightarrows NH_{4}^{+} + OH^{-}$   $0.50 - x \qquad x \qquad x$   $1.8 x 10^{-5} = \frac{(x)(x)}{(0.50 - x)} \cong \frac{x^{2}}{(0.50)}$ 

Can use the 5% rule because K < 1 and K at least 1000 time smaller than [initial]

What is the pH of a 0.50 M solution of ammonia,  $NH_3$ ,  $K_b = 1.8 \times 10^{-5}$ ?

**Step #4:** Solve for x, which is also [OH<sup>-</sup>]

 $[OH^{-}] = 3.0 \times 10^{-3} M$ 

$$\mathbf{NH}_3 + \mathbf{H}_2\mathbf{O} \leftrightarrows \mathbf{NH}_4^+ + \mathbf{OH}^-$$
  
$$\mathbf{E} \quad 0.50 - \mathbf{x} \qquad \mathbf{x} \qquad \mathbf{x}$$

$$1.8 \, x 10^{-5} = \frac{x^2}{(0.50)}$$

## <u>A Weak Base Equilibrium Problem</u>

What is the pH of a 0.50 M solution of ammonia,  $NH_3$ ,  $K_b = 1.8 \times 10^{-5}$ ?

Step #5: Convert [OH-] to pH

 $NH_3 + H_2O \leftrightarrows NH_4^+ + OH^-$ E 0.50 - x x 3.0 x 10<sup>-3</sup> M

 $pOH = -log(3.0 \times 10^{-3}) = 2.52$ pH = 14 - pOH = 14 - 2.52 = 11.48

## **Remember! Everyone always forgets!**

You can convert back and forth from Ka to Kb and vice versa. If you are given Ka for an acid but are doing problems with the acid's conjugate base you can use that Ka to find the Kb that you need.

> Ka x Kb = Kw Ka x Kb =  $(1 \times 10^{-14})$



 $Ka = \frac{[H^+][A^-]}{[HA]}$ 

## The % ionization for a weak acid **INCREASES** as concentration **DECREASES**!

Let's pretend we cut the [ ] in half

$$\mathbf{Q} = \frac{\left[\frac{1}{2}\right]\left[\frac{1}{2}\right]}{\left[\frac{1}{2}\right]} = \frac{1}{2}$$
 (of whatever the original was)

If Q decreases... Reaction shifts to make more products... More ionization! Higher % ionization!

#### YouTube Link to Presentation

https://youtu.be/WwZBnDhd9J0