

PERIODIC TABLE OF THE ELEMENTS

1 H 1.008																	2 He 4.00
3 Li 6.94	4 Be 9.01											5 B 10.81	6 C 12.01	7 N 14.01	8 O 16.00	9 F 19.00	10 Ne 20.18
11 Na 22.99	12 Mg 24.30											13 Al 26.98	14 Si 28.09	15 P 30.97	16 S 32.06	17 Cl 35.45	18 Ar 39.95
19 K 39.10	20 Ca 40.08	21 Sc 44.96	22 Ti 47.90	23 V 50.94	24 Cr 52.00	25 Mn 54.94	26 Fe 55.85	27 Co 58.93	28 Ni 58.69	29 Cu 63.55	30 Zn 65.39	31 Ga 69.72	32 Ge 72.59	33 As 74.92	34 Se 78.96	35 Br 79.90	36 Kr 83.80
37 Rb 85.47	38 Sr 87.62	39 Y 88.91	40 Zr 91.22	41 Nb 92.91	42 Mo 95.94	43 Tc (98)	44 Ru 101.1	45 Rh 102.91	46 Pd 106.42	47 Ag 107.87	48 Cd 112.41	49 In 114.82	50 Sn 118.71	51 Sb 121.75	52 Te 127.60	53 I 126.91	54 Xe 131.29
55 Cs 132.91	56 Ba 137.33	57 *La 138.91	72 Hf 178.49	73 Ta 180.95	74 W 183.85	75 Re 186.21	76 Os 190.2	77 Ir 192.2	78 Pt 195.08	79 Au 196.97	80 Hg 200.59	81 Tl 204.38	82 Pb 207.2	83 Bi 208.98	84 Po (209)	85 At (210)	86 Rn (222)
87 Fr (223)	88 Ra 226.02	89 †Ac 227.03	104 Rf (261)	105 Db (262)	106 Sg (266)	107 Bh (264)	108 Hs (277)	109 Mt (268)	110 Ds (271)	111 Rg (272)							

*Lanthanide Series

† Actinide Series

58 Ce 140.12	59 Pr 140.91	60 Nd 144.24	61 Pm (145)	62 Sm 150.4	63 Eu 151.97	64 Gd 157.25	65 Tb 158.93	66 Dy 162.50	67 Ho 164.93	68 Er 167.26	69 Tm 168.93	70 Yb 173.04	71 Lu 174.97
90 Th 232.04	91 Pa 231.04	92 U 238.03	93 Np (247)	94 Pu (244)	95 Am (243)	96 Cm (247)	97 Bk (247)	98 Cf (251)	99 Es (252)	100 Fm (257)	101 Md (258)	102 No (259)	103 Lr (262)

AP® CHEMISTRY EQUATIONS AND CONSTANTS, EFFECTIVE 2025

UNIT SYMBOLS	
gram,	g
mole,	mol
liter,	L
meter,	m
second,	s
hertz,	Hz
atmosphere,	atm
millimeter of mercury,	mm Hg
degree Celsius,	°C
Kelvin,	K
joule,	J
volt,	V
coulomb,	C
ampere,	A

UNIT CONVERSIONS
1 hertz = 1 s ⁻¹
1 atm = 760 mm Hg = 760 torr
K = °C + 273.15
1 volt = $\frac{1 \text{ joule}}{1 \text{ coulomb}}$
1 ampere = $\frac{1 \text{ coulomb}}{1 \text{ second}}$

METRIC PREFIXES		
Factor	Prefix	Symbol
10 ⁹	giga	G
10 ⁶	mega	M
10 ³	kilo	k
10 ⁻²	centi	c
10 ⁻³	milli	m
10 ⁻⁶	micro	μ
10 ⁻⁹	nano	n
10 ⁻¹²	pico	p

ATOMIC STRUCTURE

$$E = h\nu$$

$$c = \lambda\nu$$

$$F_{\text{coulombic}} \propto \frac{q_1q_2}{r^2}$$

E = energy

ν = frequency

λ = wavelength

F = force

q = charge

r = separation

Planck's constant, *h* = 6.626 × 10⁻³⁴ J s

Speed of light, *c* = 2.998 × 10⁸ m s⁻¹

Avogadro's number = 6.022 × 10²³ mol⁻¹

GASES, LIQUIDS, AND SOLUTIONS

$$\frac{P_1V_1}{T_1} = \frac{P_2V_2}{T_2}$$

$$PV = nRT$$

$$P_A = P_{\text{total}} \times X_A, \text{ where } X_A = \frac{\text{moles A}}{\text{total moles}}$$

$$P_{\text{total}} = P_A + P_B + P_C + \dots$$

$$n = \frac{m}{M}$$

$$D = \frac{m}{V}$$

$$KE = \frac{1}{2}mv^2$$

$$M = \frac{n_{\text{solute}}}{L_{\text{solution}}}$$

$$A = \epsilon bc$$

P = pressure

V = volume

T = temperature

n = number of moles

X = mole fraction

m = mass

M = molar mass

D = density

KE = kinetic energy

v = velocity

M = molarity

A = absorbance

ε = molar absorptivity

b = path length

c = concentration

Gas constant, *R* = 8.314 J mol⁻¹ K⁻¹

= 0.08206 L atm K⁻¹ mol⁻¹

STP = 273.15 K and 1.0 atm

Ideal gas at STP = 22.4 L mol⁻¹

KINETICS

$$[A]_t - [A]_0 = -kt$$

$$\ln[A]_t - \ln[A]_0 = -kt$$

$$\frac{1}{[A]_t} - \frac{1}{[A]_0} = kt$$

$$t_{1/2} = \frac{0.693}{k}$$

k = rate constant

t = time

*t*_{1/2} = half-life

EQUILIBRIUM

$$K_c = \frac{[C]^c[D]^d}{[A]^a[B]^b}, \text{ where } aA + bB \rightleftharpoons cC + dD$$

$$K_p = \frac{(P_C)^c(P_D)^d}{(P_A)^a(P_B)^b}$$

$$K_w = [H_3O^+][OH^-] = 1.0 \times 10^{-14} \text{ at } 25^\circ\text{C}$$

$$pK_w = 14 = \text{pH} + \text{pOH} \text{ at } 25^\circ\text{C}$$

$$\text{pH} = -\log[H_3O^+], \quad \text{pOH} = -\log[OH^-]$$

$$K_a = \frac{[H_3O^+][A^-]}{[HA]}, \quad K_b = \frac{[OH^-][HB^+]}{[B]}$$

$$pK_a = -\log K_a, \quad pK_b = -\log K_b$$

$$K_w = K_a \times K_b, \quad pK_w = pK_a + pK_b$$

$$\text{pH} = pK_a + \log \frac{[A^-]}{[HA]}$$

Equilibrium Constants

K_c (molar concentrations)

K_p (gas pressures)

K_w (water)

K_a (acid)

K_b (base)

THERMODYNAMICS/ELECTROCHEMISTRY

$$q = mc\Delta T$$

$$\Delta H_{\text{reaction}}^\circ = \sum \Delta H_f^\circ \text{ products} - \sum \Delta H_f^\circ \text{ reactants}$$

$$\Delta S_{\text{reaction}}^\circ = \sum S_{\text{products}}^\circ - \sum S_{\text{reactants}}^\circ$$

$$\Delta G_{\text{reaction}}^\circ = \sum \Delta G_f^\circ \text{ products} - \sum \Delta G_f^\circ \text{ reactants}$$

$$\Delta G^\circ = \Delta H^\circ - T\Delta S^\circ$$

$$= -RT \ln K$$

$$= -nFE^\circ$$

$$I = \frac{q}{t}$$

$$E_{\text{cell}} = E_{\text{cell}}^\circ - \frac{RT}{nF} \ln Q$$

q = heat

m = mass

c = specific heat capacity

T = temperature

S[°] = standard entropy

H[°] = standard enthalpy

G[°] = standard Gibbs free energy

R = gas constant

K = equilibrium constant

n = number of moles of electrons

E[°] = standard potential

I = current (amperes)

q = charge (coulombs)

t = time (seconds)

Q = reaction quotient

Faraday's constant, *F* = 96,485 coulombs / 1 mol e⁻