

6 • Energy and Chemical Reactions

PRACTICE TEST

- A system has an increase in internal energy, ΔE , of 40 kJ. If 20 kJ of work, w , is done on the system, what is the heat change, q ?
 - +60 kJ
 - +40 kJ
 - +20 kJ
 - 20 kJ
 - 60 kJ
- A gas at 20 atm pressure with a volume of 2.0 Liters expands against a 5 atm pressure to a volume of 8.0 Liters. How much work is done by the gas?
 - 30 L·atm
 - 18 L·atm
 - 8 L·atm
 - 5 L·atm
- Which equation represents the heat of formation, ΔH_f , for MgCl_2 ?
 - $\text{Mg}^{2+}(\text{aq}) + 2 \text{Cl}^- \rightarrow \text{MgCl}_2(\text{s})$
 - $\text{Mg}(\text{s}) + 2 \text{Cl}(\text{g}) \rightarrow \text{MgCl}_2(\text{s})$
 - $\text{MgCl}_2(\text{s}) \rightarrow \text{Mg}^{2+}(\text{aq}) + 2 \text{Cl}^-(\text{aq})$
 - $\text{Mg}(\text{s}) + \text{Cl}_2(\text{g}) \rightarrow \text{MgCl}_2(\text{s})$
- Take a toy balloon. Quickly stretch the balloon and press it against your lower lip. What is the ΔH for the reaction:

unstretched \rightarrow stretched

 - +
 - 0
 -
 - impossible to tell
- Which of the following is NOT a state function?
 - pressure
 - volume
 - temperature
 - none of these
- The correct units for specific heat capacity:
 - $\text{J}/^\circ\text{C}$
 - J/g
 - $\text{J}/\text{g } ^\circ\text{C}$
 - $^\circ\text{C}/\text{g}$
- How much heat is required to convert solid sulfur to gaseous sulfur at 298 K and 1 atm pressure?

	ΔH° (kJ/mol)
$\text{S}(\text{s}) + \text{O}_2(\text{g}) \rightarrow \text{SO}_2(\text{g})$	-395
$\text{S}(\text{g}) + \text{O}_2(\text{g}) \rightarrow \text{SO}_2(\text{g})$	-618

 - 1013 kJ/mol
 - 223 kJ/mol
 - +223 kJ/mol
 - +618 kJ/mol
- Using the ΔH_f° given below, calculate the $\Delta H_{\text{combustion}}$ for propane, C_3H_8 .

	ΔH_f° (kJ/mol)
$\text{H}_2\text{O}(\text{l})$	-286
$\text{CO}_2(\text{g})$	-394
$\text{C}_3\text{H}_8(\text{g})$	-104

 - 576 kJ
 - 576 kJ
 - 2222 kJ
 - 2330 kJ
- The heat of vaporization of methane, CH_4 , at its boiling point is 9.20 kJ/mol. How much heat energy is required to vaporize 100. g of methane at its boiling point?
 - 1380 kJ
 - 86.3 kJ
 - 21.6 kJ
 - 57.4 kJ
- How much energy is required to melt 10.0 g benzene, C_6H_6 ? The heat of fusion of benzene is 2.37 kJ/mol.
 - 3.30 kJ
 - 23.7 kJ
 - 1850 kJ
 - 0.303 kJ

11. If ΔH for a reaction is positive, ...
- the reaction rate is generally very fast.
 - the enthalpy change of the reverse reaction is positive.
 - the enthalpy of the products is greater than the enthalpy of the reactants.
 - the energy released during bond formation is greater than the energy absorbed during bonding breaking for the reaction.

12. Given the two equations:



What is the standard enthalpy of formation for sulfur dioxide, $\text{SO}_2(\text{g})$?

- +99.1 kJ
 - 296.1 kJ
 - 592.2 kJ
 - 839.5 kJ
13. When 0.100 g benzoic acid ($\text{HC}_6\text{H}_4\text{CO}_2$) and excess oxygen is ignited in a bomb calorimeter, the temperature of the water changes from 25.000°C to 25.225°C . The heat capacity of the calorimeter is $603 \text{ J}/^\circ\text{C}$. What is the ΔE for this reaction?
- 597 J
 - 1660 J
 - 136 J
 - 149 J
14. Under conditions of constant volume, the heat change that occurs during a chemical reaction is equal to
- ΔH
 - ΔE
 - ΔT
 - ΔP
15. Systems tend toward:
- maximum entropy and minimum enthalpy
 - maximum entropy and maximum enthalpy
 - minimum entropy and minimum enthalpy
 - minimum entropy and maximum enthalpy

Answers:

- | | | |
|------|-------|-------|
| 1. C | 6. C | 11. C |
| 2. A | 7. C | 12. B |
| 3. D | 8. C | 13. C |
| 4. C | 9. D | 14. B |
| 5. D | 10. D | 15. A |

Notes:

- an increase in internal energy means an increase in P.E. of system by 40 kJ
work done ON system increases P.E., +20 kJ, so $q = +20 \text{ kJ}$, too.
- work = $P\Delta V = 5 \text{ atm} \times (8-2 \text{ L})$... the 20 atm is not used for anything.
- balloon gets warm, $\Delta H < 0$
- each of these only depends on the STATE of the substance, not on its HISTORY.
- reverse second reaction
- recall: $\text{C}_3\text{H}_8 + 5\text{O}_2 \rightarrow 3\text{CO}_2 + 4\text{H}_2\text{O}$ and use Hess's Law
- Given: 100g CH_4 , use molar mass & H_{vap} as conversion factors.
- Given: 10.0g C_6H_6 , use molar mass & H_{fus} as conversion factors.
- this is an "uphill" reaction.
- take half of first equation, reverse second equation. if you reverse and double second equation, you get TWICE the answer.
- answer = heat capacity $\times \Delta T$... you don't use 0.100 g anywhere. You would IF the question asked for MOLAR heat of combustion.
- if volume is constant, $P\Delta V$ work = 0 so $\Delta E = q + w$ becomes $\Delta E = q$.
- from lecture... THIS chapter, however, concentrates on enthalpy, ΔH .