

2021

AP[®]

 CollegeBoard

AP[®] Chemistry

Free-Response Questions

© 2021 College Board. College Board, Advanced Placement, AP, AP Central, and the acorn logo are registered trademarks of College Board. Visit College Board on the web: collegeboard.org.

AP Central is the official online home for the AP Program: apcentral.collegeboard.org.

PERIODIC TABLE OF THE ELEMENTS

1																	18	
1																	2	
H 1.008																	He 4.00	
3																	9	
Li 6.94																	F 19.00	
11																	17	
Na 22.99																	Cl 35.45	
19																	35	
K 39.10																	Kr 83.80	
37																	53	
Rb 85.47																	Xe 131.29	
55																	85	
Cs 132.91																	Rn	
87																	117	
Fr 88																	Og 118	
	2																	17
	4																	10
Be 9.01																	Ne 20.18	
12																	18	
Mg 24.30																	Ar 39.95	
20																	36	
Ca 40.08																	Kr 83.80	
38																	54	
Sr 87.62																	Xe 131.29	
86																	86	
Ba 137.33																	Rn	
137.33																	118	
87																	118	
Ra 89-103 †																	Og 118	
	3																	12
21																	30	
Sc 44.96																	Zn 65.38	
39																	48	
Y 88.91																	Cd 112.41	
88.91																	112.41	
56																	80	
Ba 137.33																	Rn	
137.33																	118	
87																	118	
Ra 89-103 †																	Og 118	
	4																	13
22																	31	
Ti 47.87																	Ga 69.72	
40																	49	
Zr 91.22																	In 114.82	
91.22																	114.82	
72																	81	
Hf 178.49																	Tl 204.38	
178.49																	204.38	
104																	113	
Rf 104																	Nh 113	
104																	113	
88																	114	
Ra 89-103 †																	Og 118	
88																	114	
89-103 †																	115	
89-103 †																	116	
89-103 †																	117	
89-103 †																	118	
89-103 †																	119	
89-103 †																	120	
89-103 †																	121	
89-103 †																	122	
89-103 †																	123	
89-103 †																	124	
89-103 †																	125	
89-103 †																	126	
89-103 †																	127	
89-103 †																	128	
89-103 †																	129	
89-103 †																	130	
89-103 †																	131	
89-103 †																	132	
89-103 †																	133	
89-103 †																	134	
89-103 †																	135	
89-103 †																	136	
89-103 †																	137	
89-103 †																	138	
89-103 †																	139	
89-103 †																	140	
89-103 †																	141	
89-103 †																	142	
89-103 †																	143	
89-103 †																	144	
89-103 †																	145	
89-103 †																	146	
89-103 †																	147	
89-103 †																	148	
89-103 †																	149	
89-103 †																	150	
89-103 †																	151	
89-103 †																	152	
89-103 †																	153	
89-103 †																	154	
89-103 †																	155	
89-103 †																	156	
89-103 †																	157	
89-103 †																	158	
89-103 †																	159	
89-103 †																	160	
89-103 †																	161	
89-103 †																	162	
89-103 †																	163	
89-103 †																	164	
89-103 †																	165	
89-103 †																	166	
89-103 †																	167	
89-103 †																	168	
89-103 †																	169	
89-103 †																	170	
89-103 †																	171	
89-103 †																	172	
89-103 †																	173	
89-103 †																	174	
89-103 †																	175	
89-103 †																	176	
89-103 †																	177	
89-103 †																	178	
89-103 †																	179	
89-103 †																	180	
89-103 †																	181	
89-103 †																	182	
89-103 †																	183	
89-103 †																	184	
89-103 †																	185	
89-103 †																	186	
89-103 †																	187	
89-103 †																	188	
89-103 †																	189	
89-103 †																	190	
89-103 †																	191	
89-103 †																	192	
89-103 †																	193	
89-103 †																	194	
89-103 †																	195	
89-103 †																	196	
89-103 †																	197	
89-103 †																	198	
89-103 †																	199	
89-103 †																	200	
89-103 †																	201	
89-103 †																	202	
89-103 †																	203	
89-103 †																	204	
89-103 †																	205	
89-103 †																	206	
89-103 †																	207	
89-103 †																	208	
89-103 †																	209	
89-103 †																	210	
89-103 †																	211	
89-103 †																	212	
89-103 †																	213	
89-103 †																	214	
89-103 †																	215	
89-103 †																	216	
89-103 †																	217	
89-103 †																	218	
89-103 †																	219	
89-103 †																	220	
89-103 †																	221	
89-103 †																	222	
89-103 †																	223	
89-103 †																	224	
89-103 †																	225	
89-103 †																	226	
89-103 †																	227	
89-103 †																	228	
89-103 †																	229	
89-103 †																	230	
89-103 †																	231	
89-103 †																	232	
89-103 †																	233	
89-103 †																	234	
89-103 †																	235	
89-103 †																	236	
89-103 †																	237	
89-103 †																	238	
89-103 †																	239	
89-103 †																	240	
89-103 †																	241	
89-103 †																	242	
89-103 †																	243	
89-103 †																	244	
89-103 †																	245	
89-103 †																	246	
89-103 †																	247	
89-103 †																	248	
89-103 †																	249	
89-103 †																	250	
89-103 †																	251	
89-103 †																	252	
89-103 †																	253	
89-103 †																	254	
89-103 †																	255	
89-103 †																	256	
89-103 †																	257	
89-103 †																	258	
89-103 †																	259	
89-103 †																	260	
89-103 †																	261	
89-103 †																	262	
89-103 †																	263	
89-103 †																	264	
89-103 †																	265	
89-103 †																	266	
89-103 †																	267	
89-103 †																	268	
89-103 †																	269	
89-103 †																	270	
89-103 †																	271	
89-103 †																	272	
89-103 †																	273	
89-103 †																	274	
89-103 †																	275	
89-103 †																	276	
89-103 †																	277	
89-103 †																	278	
89-103 †																	279	
89-103 †																	280	
89-103 †																	281	
89-103 †																	282	
89-103 †																	283	
89-103 †																	284	
89-103 †																	285	
89-103 †																	286	
89-103 †																	287	
89-103 †																	288	
89-103 †																	289	
89-103 †																	290	
89-103 †																	291	
89-103 †																	292	
89-103 †																	293	
89-103 †																	294	
89-103 †																	295	
89-103 †																	296	
89-103 †																	297	
89-103 †																	298	
89-103 †																	299	
89-103 †																	300	
89-103 †																	301	
89-103 †																	302	
89-103 †																	303	
89-103 †																	304	
89-103 †																	305	
89-103 †																	306	
89-103 †																	307	
89-103 †																	308	
89-103 †																	309	
89-103 †																	310	
89-103 †																	311	
89-103 †																	312	
89-103 †																	313	
89-103 †																	314	
89-103 †																	315	
89-103 †																	316	
89-103 †																	317	
89-103 †																	318	
89-103 †																	319	
89-103 †																	320	
89-103 †																	321	
89-103 †																	322	
89-103 †																	323	
89-103 †																	324	
89-103 †																	325	
89-103 †																	326	
89-103 †																	327	
89-103 †																	328	
89-103 †																	329	
89-103 †																	330	
89-103 †																	331	
89-103 †																	332	
89-103 †																	333	
89-103 †																	334	
89-103 †																	335	
89-103 †																	336	
89-103 †																	337	
89-103 †																	338	
89-103 †																	339	
89-103 †																	340	
89-103 †																	341	
89-103 †																	342	
89-103 †																	343	
89-103 †																	344	
89-103 †																	345	
89-103 †																	346	
89-103 †																	347	
89-103 †																	348	
89-103 †																	349	
89-103 †																	350	
89-103 †																	351	
89-103 †																	352	
89-103 †																	353	
89-103 †																	354	
89-103 †																	355	
89-103 †																	356	
89-103 †																	357	
89-103 †																	358	
89-103 †																	359	
89-103 †																	360	
89-103 †																	361	
89-103 †																	362	
89-103 †																	363	
89-103 †																	364	

AP[®] CHEMISTRY EQUATIONS AND CONSTANTS

Throughout the exam the following symbols have the definitions specified unless otherwise noted.

L, mL = liter(s), milliliter(s)
 g = gram(s)
 nm = nanometer(s)
 atm = atmosphere(s)

mm Hg = millimeters of mercury
 J, kJ = joule(s), kilojoule(s)
 V = volt(s)
 mol = mole(s)

ATOMIC STRUCTURE

$$E = h\nu$$

$$c = \lambda\nu$$

E = energy
 ν = frequency
 λ = wavelength

Planck's constant, $h = 6.626 \times 10^{-34}$ J s

Speed of light, $c = 2.998 \times 10^8$ m s⁻¹

Avogadro's number = 6.022×10^{23} mol⁻¹

Electron charge, $e = -1.602 \times 10^{-19}$ coulomb

EQUILIBRIUM

$$K_c = \frac{[C]^c [D]^d}{[A]^a [B]^b}, \text{ where } a A + b B \rightleftharpoons c C + d D$$

$$K_p = \frac{(P_C)^c (P_D)^d}{(P_A)^a (P_B)^b}$$

$$K_a = \frac{[H^+][A^-]}{[HA]}$$

$$K_b = \frac{[OH^-][HB^+]}{[B]}$$

$$K_w = [H^+][OH^-] = 1.0 \times 10^{-14} \text{ at } 25^\circ\text{C}$$

$$= K_a \times K_b$$

$$\text{pH} = -\log[H^+], \text{ pOH} = -\log[OH^-]$$

$$14 = \text{pH} + \text{pOH}$$

$$\text{pH} = \text{p}K_a + \log \frac{[A^-]}{[HA]}$$

$$\text{p}K_a = -\log K_a, \text{ p}K_b = -\log K_b$$

Equilibrium Constants

K_c (molar concentrations)

K_p (gas pressures)

K_a (weak acid)

K_b (weak base)

K_w (water)

KINETICS

$$[A]_t - [A]_0 = -kt$$

$$\ln[A]_t - \ln[A]_0 = -kt$$

$$\frac{1}{[A]_t} - \frac{1}{[A]_0} = kt$$

$$t_{1/2} = \frac{0.693}{k}$$

k = rate constant

t = time

$t_{1/2}$ = half-life

GASES, LIQUIDS, AND SOLUTIONS

$$PV = nRT$$

$$P_A = P_{\text{total}} \times X_A, \text{ where } X_A = \frac{\text{moles A}}{\text{total moles}}$$

$$P_{\text{total}} = P_A + P_B + P_C + \dots$$

$$n = \frac{m}{M}$$

$$K = ^\circ\text{C} + 273$$

$$D = \frac{m}{V}$$

$$KE_{\text{molecule}} = \frac{1}{2}mv^2$$

Molarity, M = moles of solute per liter of solution

$$A = \varepsilon bc$$

P = pressure

V = volume

T = temperature

n = number of moles

m = mass

M = molar mass

D = density

KE = kinetic energy

v = velocity

A = absorbance

ε = molar absorptivity

b = path length

c = concentration

Gas constant, $R = 8.314 \text{ J mol}^{-1} \text{ K}^{-1}$

$= 0.08206 \text{ L atm mol}^{-1} \text{ K}^{-1}$

$= 62.36 \text{ L torr mol}^{-1} \text{ K}^{-1}$

$1 \text{ atm} = 760 \text{ mm Hg} = 760 \text{ torr}$

STP = 273.15 K and 1.0 atm

Ideal gas at STP = 22.4 L mol⁻¹

THERMODYNAMICS / ELECTROCHEMISTRY

$$q = mc\Delta T$$

$$\Delta S^\circ = \sum S^\circ \text{ products} - \sum S^\circ \text{ reactants}$$

$$\Delta H^\circ = \sum \Delta H_f^\circ \text{ products} - \sum \Delta H_f^\circ \text{ reactants}$$

$$\Delta G^\circ = \sum \Delta G_f^\circ \text{ products} - \sum \Delta G_f^\circ \text{ reactants}$$

$$\Delta G^\circ = \Delta H^\circ - T\Delta S^\circ$$

$$= -RT \ln K$$

$$= -nFE^\circ$$

$$I = \frac{q}{t}$$

$$E_{\text{cell}} = E_{\text{cell}}^\circ - \frac{RT}{nF} \ln Q$$

q = heat

m = mass

c = specific heat capacity

T = temperature

S° = standard entropy

H° = standard enthalpy

G° = standard Gibbs free energy

n = number of moles

E° = standard reduction potential

I = current (amperes)

q = charge (coulombs)

t = time (seconds)

Q = reaction quotient

Faraday's constant, $F = 96,485$ coulombs per mole of electrons


$$1 \text{ volt} = \frac{1 \text{ joule}}{1 \text{ coulomb}}$$

Begin your response to **QUESTION 1** on this page.

CHEMISTRY**SECTION II****Time—1 hour and 45 minutes****7 Questions****YOU MAY USE YOUR CALCULATOR FOR THIS SECTION.**

Directions: Questions 1–3 are long free-response questions that require about 23 minutes each to answer and are worth 10 points each. Questions 4–7 are short free-response questions that require about 9 minutes each to answer and are worth 4 points each.

For each question, show your work for each part in the space provided after that part. Examples and equations may be included in your responses where appropriate. For calculations, clearly show the method used and the steps involved in arriving at your answers. You must show your work to receive credit for your answer. Pay attention to significant figures.



1. Methanoic acid, HCOOH, ionizes according to the equation above.
 - (a) Write the expression for the equilibrium constant, K_a , for the reaction.



 - (b) Calculate the pH of a 0.25 *M* solution of HCOOH.

GO ON TO THE NEXT PAGE.

Use a pencil or pen with black or dark blue ink only. Do NOT write your name. Do NOT write outside the box.

Continue your response to **QUESTION 1** on this page.

(c) In the box below, complete the Lewis electron-dot diagram for HCOOH. Show all bonding and nonbonding valence electrons.



(d) In aqueous solution, the compound H_2NNH_2 reacts according to the equation above. A 50.0 mL sample of 0.25 M $\text{H}_2\text{NNH}_2(aq)$ is combined with a 50.0 mL sample of 0.25 M $\text{HCOOH}(aq)$.

(i) Write the balanced net ionic equation for the reaction that occurs when H_2NNH_2 is combined with HCOOH.

(ii) Is the resulting solution acidic, basic, or neutral? Justify your answer.

GO ON TO THE NEXT PAGE.


Use a pencil or pen with black or dark blue ink only. Do NOT write your name. Do NOT write outside the box.

© 2021 College Board.


Visit College Board on the web: collegeboard.org.

Continue your response to **QUESTION 1** on this page.

When a catalyst is added to a solution of $\text{HCOOH}(aq)$, the reaction represented by the following equation occurs.



(e) Is the reaction a redox reaction? Justify your answer.



(f) The reaction occurs in a rigid 4.3 L vessel at 25°C , and the total pressure is monitored, as shown in the graph above. The vessel originally did not contain any gas. Calculate the number of moles of $\text{CO}_2(g)$ produced in the reaction. (Assume that the amount of $\text{CO}_2(g)$ dissolved in the solution is negligible.)

(g) After the reaction has proceeded for several minutes, does the amount of catalyst increase, decrease, or remain the same? Justify your answer.


GO ON TO THE NEXT PAGE.

Use a pencil or pen with black or dark blue ink only. Do NOT write your name. Do NOT write outside the box.

Begin your response to **QUESTION 2** on this page.

2. Answer the following questions about the element Si and some of its compounds.

(a) The mass spectrum of a pure sample of Si is shown below.



(i) How many protons and how many neutrons are in the nucleus of an atom of the most abundant isotope of Si ?

(ii) Write the ground-state electron configuration of Si.

Two compounds that contain Si are SiO_2 and SiH_4 .

(b) At 161 K, SiH_4 boils but SiO_2 remains as a solid. Using principles of interparticle forces, explain the difference in boiling points.

GO ON TO THE NEXT PAGE.

Use a pencil or pen with black or dark blue ink only. Do NOT write your name. Do NOT write outside the box.

Continue your response to **QUESTION 2** on this page.

At high temperatures, SiH_4 decomposes to form solid silicon and hydrogen gas.

(c) Write a balanced equation for the reaction.

A table of absolute entropies of some substances is given below.

Substance	S° (J/(mol · K))
$\text{H}_2(g)$	131
$\text{Si}(s)$	18
$\text{SiH}_4(g)$	205

(d) Explain why the absolute molar entropy of $\text{Si}(s)$ is less than that of $\text{H}_2(g)$.

(e) Calculate the value, in J/(mol · K), of ΔS° for the reaction.

(f) The reaction is thermodynamically favorable at all temperatures. Explain why the reaction occurs only at high temperatures.

GO ON TO THE NEXT PAGE.


Use a pencil or pen with black or dark blue ink only. Do NOT write your name. Do NOT write outside the box.

© 2021 College Board.

Visit College Board on the web: collegeboard.org.

Continue your response to **QUESTION 2** on this page.

(g) A partial photoelectron spectrum of pure Si is shown below. On the spectrum, draw the missing peak that corresponds to the electrons in the $3p$ sublevel.



(h) Using principles of atomic structure, explain why the first ionization energy of Ge is lower than that of Si.

(i) A single photon with a wavelength of 4.00×10^{-7} m is absorbed by the Si sample. Calculate the energy of the photon in joules.

GO ON TO THE NEXT PAGE.

Use a pencil or pen with black or dark blue ink only. Do NOT write your name. Do NOT write outside the box.

Begin your response to **QUESTION 3** on this page.

3. A student is given the task of determining the molar concentration of a CuSO_4 solution using two different procedures, precipitation and spectrophotometry.

For the precipitation experiment, the student adds 20.0 mL of 0.200 M $\text{Ba}(\text{NO}_3)_2$ to 50.0 mL of the $\text{CuSO}_4(aq)$. The reaction goes to completion, and a white precipitate forms. The student filters the precipitate and dries it overnight. The data are given in the following table.

Mass of dry filter paper	0.764 g
Volume of $\text{CuSO}_4(aq)$	50.0 mL
Volume of 0.200 M $\text{Ba}(\text{NO}_3)_2$	20.0 mL
Mass of filter paper and dried precipitate	1.136 g

(a) Write a balanced net ionic equation for the precipitation reaction.

(b) Calculate the number of moles of precipitate formed.

(c) Calculate the molarity of the original CuSO_4 solution.

GO ON TO THE NEXT PAGE.

Use a pencil or pen with black or dark blue ink only. Do NOT write your name. Do NOT write outside the box.

© 2021 College Board.

Visit College Board on the web: collegeboard.org.

Continue your response to **QUESTION 3** on this page.

For the spectrophotometry experiment, the student first makes a standard curve. The student uses a 0.1000 *M* solution of $\text{CuSO}_4(aq)$ to make three more solutions of known concentration (0.0500 *M*, 0.0300 *M*, and 0.0100 *M*) in 50.00 mL volumetric flasks.

(d) Calculate the volume of 0.1000 *M* $\text{CuSO}_4(aq)$ needed to make 50.00 mL of 0.0500 *M* $\text{CuSO}_4(aq)$.


(e) Briefly describe the procedure the student should follow to make 50.00 mL of 0.0500 *M* $\text{CuSO}_4(aq)$ using 0.1000 *M* $\text{CuSO}_4(aq)$, a 50.00 mL volumetric flask, and other standard laboratory equipment. Assume that all appropriate safety precautions will be taken.

GO ON TO THE NEXT PAGE.

Use a pencil or pen with black or dark blue ink only. Do NOT write your name. Do NOT write outside the box.

Continue your response to **QUESTION 3** on this page.

The standard curve is given below.



(f) The absorbance of the CuSO_4 solution of unknown concentration is 0.219. Determine the molarity of the solution.

(g) A second student performs the same experiment. There are a few drops of water in the cuvette before the second student adds the $\text{CuSO}_4(aq)$ solution of unknown concentration. Will this result in a $\text{CuSO}_4(aq)$ concentration for the unknown that is greater than, less than, or equal to the concentration determined in part (f)? Justify your answer.


GO ON TO THE NEXT PAGE.

Use a pencil or pen with black or dark blue ink only. Do NOT write your name. Do NOT write outside the box.

© 2021 College Board.

Visit College Board on the web: collegeboard.org.

Begin your response to **QUESTION 4** on this page.



4. A student investigates a reaction used in hand warmers, represented above. The student mixes Fe(s) with a catalyst and sand in a small open container. The student measures the temperature of the mixture as the reaction proceeds. The data are given in the following table.

Time (min)	Temperature of Mixture (°C)
0	22.0
1	25.1
2	34.6
3	37.3
4	39.7
5	39.4

(a) The mixture (Fe(s), catalyst, and sand) has a total mass of 15.0 g and a specific heat capacity of 0.72 J/(g·°C). Calculate the amount of heat absorbed by the mixture from 0 minutes to 4 minutes.

(b) Calculate the mass of Fe(s), in grams, that reacted to generate the amount of heat calculated in part (a).

GO ON TO THE NEXT PAGE.

Use a pencil or pen with black or dark blue ink only. Do NOT write your name. Do NOT write outside the box.

Continue your response to **QUESTION 4** on this page.

(c) In a second experiment, the student uses twice the mass of iron as that calculated in part (b) but the same mass of sand as in the first experiment. Would the maximum temperature reached in the second experiment be greater than, less than, or equal to the maximum temperature in the first experiment? Justify your answer.


GO ON TO THE NEXT PAGE.

Use a pencil or pen with black or dark blue ink only. Do NOT write your name. Do NOT write outside the box.

© 2021 College Board.

Visit College Board on the web: collegeboard.org.

Begin your response to **QUESTION 5** on this page.



Half-Reaction	E° (V)
$\text{Mg}^{2+} + 2 e^- \rightarrow \text{Mg}$	-2.37
$\text{Cl}_2 + 2 e^- \rightarrow 2 \text{Cl}^-$	+1.36

5. Molten MgCl_2 can be decomposed into its elements if a sufficient voltage is applied using inert electrodes. The products of the reaction are liquid Mg (at the cathode) and Cl_2 gas (at the anode). A simplified representation of the cell is shown above. The reduction half-reactions related to the overall reaction in the cell are given in the table.

(a) Draw an arrow on the diagram to show the direction of electron flow through the external circuit as the cell operates.

(b) Would an applied voltage of 2.0 V be sufficient for the reaction to occur? Support your claim with a calculation as part of your answer.

(c) If the current in the cell is kept at a constant 5.00 amps, how many seconds does it take to produce 2.00 g of $\text{Mg}(l)$ at the cathode?

GO ON TO THE NEXT PAGE.

Use a pencil or pen with black or dark blue ink only. Do NOT write your name. Do NOT write outside the box.

Begin your response to **QUESTION 6** on this page.

6. A student is studying the properties of CaSO_4 and PbSO_4 . The student has samples of both compounds, which are white powders.
- (a) The student tests the electrical conductivity of each solid and observes that neither solid conducts electricity. Describe the structures of the solids that account for their inability to conduct electricity.

The student places excess $\text{CaSO}_4(s)$ in a beaker containing 100 mL of water and places excess $\text{PbSO}_4(s)$ in another beaker containing 100 mL of water. The student stirs the contents of the beakers and then measures the electrical conductivity of the solution in each beaker. The student observes that the conductivity of the solution in the beaker containing the $\text{CaSO}_4(s)$ is higher than the conductivity of the solution in the beaker containing the $\text{PbSO}_4(s)$.

- (b) Which compound is more soluble in water, $\text{CaSO}_4(s)$ or $\text{PbSO}_4(s)$? Justify your answer based on the results of the conductivity test.

GO ON TO THE NEXT PAGE.


Use a pencil or pen with black or dark blue ink only. Do NOT write your name. Do NOT write outside the box.

© 2021 College Board.

Visit College Board on the web: collegeboard.org.

Continue your response to **QUESTION 6** on this page.

The left side of the diagram below shows a particulate representation of the contents of the beaker containing the $\text{CaSO}_4(s)$ from the solution conductivity experiment.




(c) Draw a particulate representation of $\text{PbSO}_4(s)$ and the ions dissolved in the solution in the beaker on the right in the diagram. Draw the particles to look like those shown to the right of the beaker. Draw an appropriate number of dissolved ions relative to the number of dissolved ions in the beaker on the left.

(d) The student attempts to increase the solubility of $\text{CaSO}_4(s)$ by adding 10.0 mL of 2 M $\text{H}_2\text{SO}_4(aq)$ to the beaker, and observes that additional precipitate forms in the beaker. Explain this observation.

GO ON TO THE NEXT PAGE.

Use a pencil or pen with black or dark blue ink only. Do NOT write your name. Do NOT write outside the box.

Begin your response to **QUESTION 7** on this page.



7. A student investigates gas behavior using a rigid cylinder with a movable piston of negligible mass, as shown in the diagram above. The cylinder contains 0.325 mol of O₂(g).

(a) The cylinder has a volume of 7.95 L at 25°C and 1.00 atm. Calculate the density of the O₂(g), in g/L, under these conditions.

(b) Attempting to change the density of the O₂(g), the student opens the valve on the side of the cylinder, pushes down on the piston to release some of the gas, and closes the valve again. The temperature of the gas remains constant at 25°C. Will this action change the density of the gas remaining in the cylinder? Justify your answer.

GO ON TO THE NEXT PAGE.

Use a pencil or pen with black or dark blue ink only. Do NOT write your name. Do NOT write outside the box.

© 2021 College Board.

Visit College Board on the web: collegeboard.org.

Continue your response to **QUESTION 7** on this page.

(c) The student tries to change the density of the $\text{O}_2(g)$ by cooling the cylinder to -55°C , which causes the volume of the gas to decrease. Using principles of kinetic molecular theory, explain why the volume of the $\text{O}_2(g)$ decreases when the temperature decreases to -55°C .

(d) The student further cools the cylinder to -180°C and observes that the measured volume of the $\text{O}_2(g)$ is substantially smaller than the volume that is calculated using the ideal gas law. Assume all equipment is functioning properly. Explain why the measured volume of the $\text{O}_2(g)$ is smaller than the calculated volume. (The boiling point of $\text{O}_2(l)$ is -183°C .)

GO ON TO THE NEXT PAGE.

Use a pencil or pen with black or dark blue ink only. Do NOT write your name. Do NOT write outside the box.

STOP

END OF EXAM