

Name:

Date:

Period:

Seat #:

Show all work. Complete the following on binder paper and BOX your final answers.

[1] Consider the decomposition of H_2O_2 (hydrogen peroxide) at 298 K and 1 atm pressure according to:



Substance	$\Delta G_f^\circ \left(\frac{\text{kJ}}{\text{mol}} \right)$	$\Delta H_f^\circ \left(\frac{\text{kJ}}{\text{mol}} \right)$	$S^\circ (\text{J/mol}^{-1}\text{K}^{-1})$
$\text{H}_2\text{O}_2 (l)$	-120.2	-187.6	109.5
$\text{H}_2\text{O} (l)$	-237.0	-285.8	69.4
$\text{O}_2 (g)$	----	----	205

Please find the

a) standard enthalpy of reaction


b) standard entropy of reaction.

[2] For the reaction in the previous question, please find the


a) standard (Gibbs) free energy of reaction

b) the value of the (thermodynamic) equilibrium constant at 298 K, 1 atm

[3] From the indicated standard enthalpies of formation given in kJ/mol, please calculate the standard enthalpy of reaction for:




[4] Carbon monoxide in the atmosphere slowly converts to carbon dioxide at normal atmospheric temperatures according to:




The standard enthalpy of reaction is -284 kJ and the standard entropy of reaction is -87 J/K. Estimate the temperature at which the equilibrium begins to favor the decomposition of CO_2 . Assume that the enthalpy and the entropy of reaction are not affected by temperature

[5] From the indicated standard entropies given in J/K, please calculate the standard entropy of reaction for:



192.5 121.2 130.6


[6] Please calculate the standard (Gibbs) free energy of reaction for:



Substance	$\Delta G_f^\circ \left(\frac{\text{kJ}}{\text{mol}} \right)$	$\Delta H_f^\circ \left(\frac{\text{kJ}}{\text{mol}} \right)$	$S^\circ (\text{J/mol}^{-1}\text{K}^{-1})$
NO (g)	86.69	90.4	210.6
NO ₂ (g)	51.84	34.0	240.5
O ₂ (g)	----	----	205.0

[7] Calculate the entropy of vaporization of propane given that its enthalpy of vaporization is 16.9 kJ/mol at its normal boiling point of -42.1°C.

[8] Obtain the numerical value of the equilibrium constant (at 298K) for the following reaction:



Substance	$\Delta G_f^\circ \left(\frac{\text{kJ}}{\text{mol}} \right)$	$\Delta H_f^\circ \left(\frac{\text{kJ}}{\text{mol}} \right)$	$S^\circ (\text{J/mol}^{-1}\text{K}^{-1})$
H ₂ CO ₃ (aq)	-623	-700	187
H ₂ O(liq)	-237	-286	70
CO ₂ (gas)	-394	-394	213

[9] Please indicate if TRUE or FALSE (Explain why as well):

	The entropy of a gas increases with increasing temperature
	The energy of a perfect crystal is zero at 0 K.
	Spontaneous processes always increase the entropy of the reacting system
	All spontaneous processes release heat to the surroundings
	An endothermic reaction is more likely to be spontaneous at high temperatures than at low temperatures
	The entropy of sugar decreases as it precipitates from an aqueous solution

[10] Ammonia gas a standard (Gibbs) free energy of formation equal to -16.367 kJ/mol

a) Find ΔG° for the reaction: $\text{N}_2 (\text{g}) + 3 \text{H}_2 (\text{g}) \rightleftharpoons 2 \text{NH}_3 (\text{g})$
b) In which direction will this reaction proceed if a mixture of gases is made with: $P_{\text{NH}_3} = 1.00 \text{ atm}$ $P_{\text{H}_2} = 0.50 \text{ atm}$ $P_{\text{N}_2} = 0.50 \text{ atm}$
c) What pressure of hydrogen gas should be added to a mixture already containing 0.20 atm NH_3 and 0.50 atm N_2 if one does not want the amounts of NH_3 and N_2 to change?

ANSWERS:

1. -196.4 kJ; 125 J/K
2. -233.6 kJ; 8.85×10^4
3. 175.9 kJ
4. 3260 K or 3300K
5. -133.2 J/K
6. -69.7 kJ
7. 73.1 J/K
8. 4.0×10^{-2} (using ΔG_f° data); 3.1×10^{-2} (using ΔH_f° and S° data)
9. T F F F T T
10. -32.734 kJ; proceed to the right; $5.3 \times 10^{-3} \text{ atm}$