


Name: _____

Period: _____

Seat#: _____

Directions: Complete the following chart by choosing from the following options:

Equilibrium Shift: *left, right, no change* $\Delta []$: *increase, decrease, no change* ΔK_{eq} : *no, yes*



Stressor	Equilibrium Shift	$\Delta [\text{N}_2]$	$\Delta [\text{H}_2]$	$\Delta [\text{NH}_3]$	ΔTemp	ΔK_{eq}
1) Add N ₂	Right	<i>Slight</i> increase	Decrease	Increase	Increase	No
2) Add H ₂						
3) Add NH ₃						
4) Remove N ₂						
5) Remove H ₂						
6) Remove NH ₃						
7) Increase Temp						
8) Decrease Temp						
9) Increase Pressure						
10) Decrease Pressure						
11) Write the equilibrium constant expression for K_{eq} $\text{POCl}_3(\text{g}) \leftrightarrow \text{POCl}(\text{g}) + \text{Cl}_2(\text{g})$				12) Write the equilibrium constant expression for K_{eq} $2\text{H}_2(\text{g}) + \text{O}_2(\text{g}) \leftrightarrow 2\text{H}_2\text{O}(\text{g})$		
13) Write the equilibrium constant expression for K_{eq} $2\text{C}_2\text{H}_4(\text{g}) + \text{O}_2(\text{g}) \leftrightarrow 2\text{CH}_3\text{CHO}(\text{g})$				14) Write the equilibrium constant expression for K_{eq} $2\text{H}_2\text{S}(\text{g}) + 3\text{O}_2(\text{g}) \leftrightarrow 2\text{H}_2\text{O}(\text{g}) + 2\text{SO}_2(\text{g})$		

Dougherty Valley HS Chemistry
Equilibrium – Le Chatelier’s Principle and Keq Practice

15) Write the equilibrium constant K_c for the equation $2\text{SO}_2(\text{g}) + \text{O}_2(\text{g}) \leftrightarrow 2\text{SO}_3(\text{g})$	16) Write the equilibrium constant K_c for the equation $\text{CaCO}_3(\text{s}) \leftrightarrow \text{CaO}(\text{s}) + \text{O}_2(\text{g})$
17) Write the equilibrium constant K_p for the equation $2\text{SO}_2(\text{g}) + \text{O}_2(\text{g}) \leftrightarrow 2\text{SO}_3(\text{g})$	18) Write the equilibrium constant K_p for the equation $\text{H}_2\text{O}(\text{g}) + \text{C}(\text{s}) \leftrightarrow \text{H}_2(\text{g}) + \text{CO}(\text{g})$
19) The equilibrium constant expression for a gas reaction in $K_{\text{eq}} = \frac{[\text{NH}_3]^4[\text{O}_2]^5}{[\text{NO}]^4[\text{H}_2\text{O}]^6}$ Write the balanced chemical equation corresponding to this expression.	
20) The equilibrium constant expression for a gas reaction in $K_{\text{eq}} = \frac{[\text{CS}_2][\text{H}_2]^4}{[\text{CH}_4][\text{H}_2\text{S}]^2}$ Write the balanced chemical equation corresponding to this expression.	
21) The equilibrium constant K_{eq} for the equation $2\text{HI}(\text{g}) \leftrightarrow \text{H}_2(\text{g}) + \text{I}_2(\text{g})$ at 425°C is 1.84. What is the value of K_{eq} for the following equation: $\text{H}_2(\text{g}) + \text{I}_2(\text{g}) \leftrightarrow 2\text{HI}(\text{g})$	
22) Consider the decomposition of nitrous oxide, also known as laughing gas. $2\text{N}_2\text{O}(\text{g}) \leftrightarrow 2\text{N}_2(\text{g}) + \text{O}_2(\text{g})$ At 25°C the K_c is 7.3×10^{34} a. Based on the information given, what can you say about the rate of decomposition of the reaction? b. Based on the information given, does nitrous oxide have a tendency to decompose into nitrogen and oxygen, or does it have a tendency to stay as nitrous oxide? Justify your answer. c. You can convert back and forth between K_c and K_p if you are given one of the values using the following equation: $K_p = K_c(\text{RT})^{\Delta n}$ where R is the ideal gas constant ($0.0821 \text{ L}\cdot\text{atm}/\text{K}\cdot\text{mol}$), T is temperature (in Kelvin), and Δn is the change in number of moles of gaseous products compared to gaseous reactants $\Delta n = \Sigma(\text{moles of gaseous products}) - \Sigma(\text{moles of gaseous reactants})$ Using this information, and the information given at the start of the problem, calculate the value of K_p	

Dougherty Valley HS Chemistry
Equilibrium – Le Chatelier’s Principle and Keq Practice

23) For the equilibrium system described by $2\text{SO}_2(\text{g}) + \text{O}_2(\text{g}) \leftrightarrow 2\text{SO}_3(\text{g})$ at a particular temperature the equilibrium concentrations of SO_2 , O_2 and SO_3 were 0.75 M, 0.30 M, and 0.15 M, respectively. At the temperature of the equilibrium mixture, calculate the equilibrium constant K_{eq} for the reaction.

24) For the equilibrium system described by: $\text{PCl}_5(\text{g}) \leftrightarrow \text{PCl}_3(\text{g}) + \text{Cl}_2(\text{g})$ K_{eq} equals 35 at 487°C . If the concentrations of the PCl_5 and PCl_3 are 0.015 M and 0.78 M, respectively, what is the concentration of the Cl_2 ?

25) $\text{CO}_2(\text{g}) + \text{H}_2(\text{g}) \leftrightarrow \text{CO}(\text{g}) + \text{H}_2\text{O}(\text{g})$

Calculate the value of the equilibrium constant, K_c , for the above system, if 0.1908 moles of CO_2 , 0.0908 moles of H_2 , 0.0092 moles of CO , and 0.0092 moles of H_2O vapor were present in a 2.00 L reaction vessel at equilibrium.
(Remember that $M = \text{mol/L}$)

26) The following table gives some values for reactant and product equilibrium concentrations (in mol/L) at 700 K for the Shift Reaction, an important method for the commercial production of hydrogen gas. $\text{CO}(\text{g}) + \text{H}_2\text{O}(\text{g}) \leftrightarrow \text{CO}_2(\text{g}) + \text{H}_2(\text{g})$

Trial	$[\text{CO}_2]$	$[\text{H}_2]$	$[\text{CO}]$	$[\text{H}_2\text{O}]$
1	0.600	0.600	0.266	0.266
2	0.600	0.800	0.330	0.286
3	2.00	2.00	0.877	0.896
4	1.00	1.50	0.450	0.655
5	1.80	2.00	0.590	1.20

a. Write the expression for calculating K_{eq} for the reaction.

b. Calculate the K_{eq} for each of the five trials.

c. How do the K_{eq} s for each trial compare to each other? Why?