

**Spring Final Exam Practice Test 3****- KEY****PRACTICE TEST**

The following information may be helpful.

$D = \frac{m}{V}$	$K = ^\circ\text{C} + 273$	$\underline{M} = \frac{n}{V}$	$V_c \cdot \underline{M}_c = V_d \cdot \underline{M}_d$
$\frac{P_1 V_1}{T_1} = \frac{P_2 V_2}{T_2}$	$P \cdot V = n \cdot R \cdot T$	$R = 0.0821 \frac{\text{L} \cdot \text{atm}}{\text{mol} \cdot \text{K}}$	Molar Mass = $\frac{m}{n}$
$[\text{H}^+] \cdot [\text{OH}^-] = 1.0 \times 10^{-14}$	$\text{pH} + \text{pOH} = 14$	$\text{pH} = -\log[\text{H}^+]$	$\text{pOH} = -\log[\text{OH}^-]$
$A = a \cdot b \cdot c$	$\Delta H_{\text{rxn}} = \frac{q_{\text{rxn}}}{n_{\text{rxn}}}$	$[\text{H}^+] = 10^{-\text{pH}}$	$[\text{OH}^-] = 10^{-\text{pOH}}$
1 atm = 760 mmHg = 760 torr = 14.7 psi = 101.3 kPa      1 mole = $6.022 \times 10^{23}$ particles = 22.4 L gas at STP			

**Solubility Rules for Salts**Always soluble:• alkali ions,  $\text{NH}_4^+$ ,  $\text{NO}_3^-$ ,  $\text{ClO}_3^-$ ,  $\text{ClO}_4^-$ ,  $\text{C}_2\text{H}_3\text{O}_2^-$ ,  $\text{HCO}_3^-$ Generally soluble:

•  $\text{Cl}^-$ ,  $\text{Br}^-$ ,  $\text{I}^-$  Soluble except with  $\text{Ag}^+$ ,  $\text{Pb}^{2+}$ ,  $\text{Hg}_2^{2+}$   
 •  $\text{F}^-$  Soluble except with  $\text{Pb}^{2+}$ ,  $\text{Ca}^{2+}$ ,  $\text{Ba}^{2+}$ ,  $\text{Sr}^{2+}$ ,  $\text{Mg}^{2+}$   
 •  $\text{SO}_4^{2-}$  Soluble except with  $\text{Pb}^{2+}$ ,  $\text{Ca}^{2+}$ ,  $\text{Ba}^{2+}$ ,  $\text{Sr}^{2+}$

Generally insoluble:

•  $\text{O}^{2-}$ ,  $\text{OH}^-$  Insoluble except with  $\text{Ca}^{2+}$ ,  $\text{Ba}^{2+}$ ,  $\text{Sr}^{2+}$ , alkali ions,  $\text{NH}_4^+$   
 •  $\text{CO}_3^{2-}$ ,  $\text{PO}_4^{3-}$ ,  $\text{S}^{2-}$ ,  $\text{SO}_3^{2-}$ ,  $\text{CrO}_4^{2-}$ ,  $\text{C}_2\text{O}_4^{2-}$  Insoluble except with alkali ions,  $\text{NH}_4^+$

**Strong Acids**HCl, HBr, HI,  $\text{HNO}_3$ ,  $\text{H}_2\text{SO}_4$ ,  $\text{HClO}_3$ ,  $\text{HClO}_4$ ,  $\text{HIO}_4$ **Gases that Form**

$\rightarrow \text{H}_2\text{S}(\text{g})$   $\rightarrow \text{H}_2\text{SO}_3(\text{aq}) \rightarrow \text{SO}_2(\text{g}) + \text{H}_2\text{O}(\ell)$   
 $\rightarrow \text{H}_2\text{CO}_3(\text{aq}) \rightarrow \text{CO}_2(\text{g}) + \text{H}_2\text{O}(\ell)$   $\rightarrow \text{NH}_4\text{OH}(\text{aq}) \rightarrow \text{NH}_3(\text{g}) + \text{H}_2\text{O}(\ell)$

**Part 1: Multiple Choice.** Select the answer choice that best completes the question.

- Bonds between which of the following elements is expected to be most polar?  
(A) C–H  
(B) N–O  
(C) **P–F**  
(D) S–Br
- Which type of bond is described as “a lattice of positive ions in a sea of electrons”?  
(A) covalent  
(B) hydrogen  
(C) ionic  
(D) **metallic**
- The compound  $(\text{NH}_4)_2\text{CO}_3$  contains how many atoms of hydrogen?  
(A) 2  
(B) 4  
(C) 6  
(D) **8**
- An unknown element X forms a salt with the formula  $\text{XO}_2$ . Which of the following could be X?  
(A)  $\text{Al}^{3+}$   
(B)  $\text{Ca}^{2+}$   
(C)  $\text{Na}^+$   
(D)  **$\text{Sn}^{4+}$**
- Which of the following name is correctly paired with its formula?  
(A) **Carbonate**  $\text{CO}_3^{2-}$   
(B) Chloride  $\text{ClO}_3^-$   
(C) Ferric  $\text{Fe}^{2+}$   
(D) Nitrate  $\text{NO}_2^-$
- Which one of the following is the correct formula for calcium phosphate?  
(A)  $\text{Ca}_2(\text{PO}_4)_3$   
(B)  **$\text{Ca}_3(\text{PO}_4)_2$**   
(C)  $\text{CaPO}_4$   
(D)  $\text{PO}_4\text{Ca}_3$

7. What is the name of the compound  $\text{N}_2\text{O}_3$ ?  
 (A) dinitride trioxide  
**(B) dinitrogen trioxide**  
 (C) nitrate  
 (D) nitrogen oxide
8. What is the percent of carbon in barium carbonate,  $\text{BaCO}_3$  (molar mass = 197.3 g/mol)?  
 (A) 3.04%  
**(B) 6.09%**  
 (C) 14.0%  
 (D) 20.0%
9. How many moles of hydrogen cyanide, HCN, are contained in 9.00 grams of HCN? (molar mass = 27.03 g/mol)  
**(A) 0.333**  
 (B) 0.900  
 (C) 1.00  
 (D) 9.00
10. How many molecules are in  $2.00 \times 10^{-2}$  moles of carbon tetrachloride,  $\text{CCl}_4$ ? (molar mass = 154 g/mol)  
 (A)  $1.20 \times 10^{23}$   
 (B)  $3.01 \times 10^{23}$   
 (C)  $6.02 \times 10^{23}$   
**(D)  $1.20 \times 10^{22}$**
11. Which of the following describes the bonds in a molecule of ethyne,  $\text{C}_2\text{H}_2$ ?  
 (A) 1 double bond, 2 single bonds.  
**(B) 1 triple bond, 2 single bonds.**  
 (C) 2 double bonds, 1 single bond.  
 (D) 3 single bonds.
12. The total number of dots drawn in the Lewis structure of nitrogen,  $\text{N}_2$ , is  
 (A) 5  
**(B) 10**  
 (C) 14  
 (D) 16
13. Which of the following is the correct Lewis Structure for carbon monoxide, CO?  
 (A)  $\text{:}\ddot{\text{C}}\text{:}\ddot{\text{O}}\text{:}$   
 (B)  $\text{:}\ddot{\text{C}}\text{:}\ddot{\text{O}}\text{:}$   
 (C)  $\text{:}\dot{\text{C}}\text{:}\ddot{\text{O}}\text{:}$   
**(D)  $\text{:}\text{C}\text{:}\text{O}\text{:}$**
14. Which one of these molecules is polar?  
 (A)  $\text{:}\ddot{\text{Cl}}\text{:}\ddot{\text{Cl}}\text{:}$   
**(B)**  
 $\begin{array}{c} \text{H} \\ | \\ \text{H} \text{ : } \ddot{\text{N}} \text{ : } \text{H} \end{array}$   
 (C)  $\text{:}\ddot{\text{O}}\text{:}\text{C}\text{:}\ddot{\text{O}}\text{:}$   
 (D)  
 $\begin{array}{c} \text{:}\ddot{\text{Cl}}\text{:} \\ | \\ \text{:}\ddot{\text{Cl}}\text{:}\text{C}\text{:}\ddot{\text{Cl}}\text{:} \\ | \\ \text{:}\ddot{\text{Cl}}\text{:} \end{array}$
15. Which of the following is paired **INCORRECTLY** with the solid it forms?  
 (A) Bronze – Metallic solid  
 (B) CaO – Ionic solid  
 (C)  $\text{H}_2\text{O}$  – Molecular solid  
**(D)  $\text{P}_4\text{O}_{10}$  – Network covalent solid**
16. An unknown solid has a melting point of  $400^\circ\text{C}$  and conducts electricity. Which of the following could be the solid?  
 (A) Diamond, C  
 (B) Salt, NaCl  
 (C) Sugar,  $\text{C}_6\text{H}_{12}\text{O}_6$   
**(D) Zinc, Zn**
17. Which of the following is expected for a sample of solid  $\text{P}_4\text{O}_{10}$ ?  
 (A) It is malleable.  
**(B) It has a low melting point.**  
 (C) It conducts electricity when it is melted.  
 (D) It is lustrous.
18. What intermolecular forces are present between molecules of HCl but not  $\text{H}_2$ ?  
**(A) Dipole-dipole interactions**  
 (B) Hydrogen bonding  
 (C) Ionic bonds  
 (D) London dispersion forces

19. Water molecules are attracted to each other with this type of intermolecular force.

- (A) Covalent bonding
- (B) Dipole-dipole interactions
- (C) **Hydrogen bonding**
- (D) London dispersion forces


20. Which of the following is expected to have the highest melting point?

- (A)  $\text{BH}_3$
- (B)  $\text{CH}_4$
- (C)  $\text{H}_2$
- (D)  **$\text{NH}_3$**

21. The general formula  $\text{C}_n\text{H}_{2n}$  describes the molecular composition of the hydrocarbon family known as the

- (A) alkadienes
- (B) alkanes
- (C) **alkenes**
- (D) alkynes

22. (SKIP) How many carbons make up the "parent chain" in the following molecule?




- (A) 4
- (B) 5
- (C) **6**
- (D) 7

23. (SKIP) The number of isomers of bromopropane,  $\text{C}_3\text{H}_7\text{Br}$ , is


- (A) **2**
- (B) 3
- (C) 4
- (D) 7

24. (SKIP) Which of the following is the correct name for the following compound:




- (A) **1,1-dibromobutane**
- (B) 1,1-dibromopropane
- (C) 2-bromopropane
- (D) 3,3-dibromopropane

25. (SKIP) Which of the following is an isomer of the compound below:




- (A)
- (B)
- (C)
- (D)

26. (SKIP) Which functional group is found in the following compound?




- (A) Aldehyde
- (B) Carboxylic Acid
- (C) Ether
- (D) **Ketone**

27. (SKIP) The structure of which compound is given below?



- (A) Grain alcohol (ethanol)  
 (B) Isopropyl alcohol (2-propanol)  
 (C) Nail polish remover (acetone, or propanone)  
 (D) Vinegar (ethanoic acid)

28. Which set of coefficients balances the equation for the complete combustion of ethane,  $\text{C}_2\text{H}_6$ ?



- (A) 1,3,2,3  
 (B) 1,6,2,6  
 (C) 2,6,4,5  
 (D) **2,7,4,6**

29. When the equation for the combustion of propene,  $\text{C}_3\text{H}_6$ , is balanced with the lowest whole-number coefficients, what is the coefficient of oxygen,  $\text{O}_2$ ?

- (A) 6  
 (B) **9**  
 (C) 12  
 (D) 18

30. (SKIP) When an alcohol reacts with a carboxylic acid, which of the following is formed?

- (A) A ketone  
 (B) **An ester**  
 (C) An ether  
 (D)  $\text{CO}_2$

31. Which of the following compounds is expected to be soluble in water?

- (A)  $\text{CaSO}_4$   
 (B)  $\text{FeS}$   
 (C)  $\text{PbCl}_2$   
 (D)  **$\text{SrBr}_2$**

32. Which of the following represents the dissociation of  $\text{CaI}_2$  in solution?

- (A)  $\text{CaI}_2 \rightarrow \text{Ca} + \text{I}_2$   
 (B)  **$\text{CaI}_2 \rightarrow \text{Ca}^{2+} + 2 \text{I}^-$**   
 (C)  $\text{CaI}_2 \rightarrow \text{Ca}^{2+} + \text{I}_2$   
 (D)  $\text{CaI}_2$  is insoluble so it does not dissociate.

33. How many grams of sodium hydroxide pellets,  $\text{NaOH}$ , are required to prepare 50.0 mL of a 0.150 M solution? [molar mass  $\text{NaOH} = 40.0 \text{ g/mol}$ ]

- (A) **0.300**  
 (B) 2.00  
 (C) 3.00  
 (D) 200.

34. List the following solutions prepared with the same solute in order of increasing concentration:

- I. 30.0 g solute in a 240 mL solution  
 II. 30.0 g solute in a 120 mL solution  
 III. 60.0 g solute in a 120 mL solution

- (A) **I < II < III**  
 (B) II < I < III  
 (C) II < III < I  
 (D) III < II < I

35. A 100 mL sample of a solution with a concentration of 5.00 M is diluted to a new volume of 400 mL with distilled water. The new concentration will be

- (A) **1.25 M**  
 (B) 1.66 M  
 (C) 15.0 M  
 (D) 20.0 M

36. The acid  $\text{HClO}_3$  is named:

- (A) **Chloric acid**  
 (B) Hydrochloric acid  
 (C) Hydrogen chlorate  
 (D) Hydrogen chlorine trioxide


37. A solution tested with cabbage juice turns green. Which of the following could be its pH?

- (A) 1  
 (B) 4  
 (C) **7**  
 (D) 10

38. A substance that turns cabbage juice blue and slightly lights up a light bulb is a:

- (A) strong acid  
 (B) strong base  
 (C) weak acid  
 (D) **weak base**

39. Which chemical is the conjugate base in the reaction?




- (A)  $\text{H}_2\text{O}$   
(B)  $\text{NH}_3$   
(C)  $\text{NH}_4^+$   
(D)  $\text{OH}^-$

40. If a solution has  $[\text{OH}^-] = 1.0 \times 10^{-3} \text{ M}$ , what is the pH?

- (A) 3  
(B) 7  
(C) 11  
(D) 14

41. An acid was neutralized by the following reaction:



This reaction would be classified as

- (A) decomposition  
(B) **double replacement**  
(C) single replacement  
(D) synthesis

42.  $\dots \text{H}_3\text{PO}_3 \rightarrow \dots \text{H}_3\text{PO}_4 + \dots \text{PH}_3$

When the equation above is balanced, the coefficient for  $\text{H}_3\text{PO}_4$  is:

- (A) 1  
(B) 2  
(C) 3  
(D) 4

43. When solutions of potassium sulfate and calcium bromide are combined, which of the following precipitates?

- (A)  $\text{KBr}$   
(B)  $\text{CaS}$   
(C)  **$\text{CaSO}_4$**   
(D) There is no precipitate.

44. Which of the following are products when magnesium metal is placed in hydrochloric acid?

- (A)  $\text{H}$   
(B)  $\text{H}^+$   
(C)  $\text{Mg}$   
(D)  **$\text{MgCl}_2$**

45.  $\text{Zn} \rightarrow \text{Zn}^{2+} + 2\text{e}^-$

(SKIP) In the half reaction above:

- (A) Zn is oxidized because it is gaining electrons.  
(B) **Zn is oxidized because it is losing electrons.**  
(C) Zn is reduced because it is gaining electrons.  
(D) Zn is reduced because it is losing electrons.

46. (SKIP) The classic holiday demonstration, "Chemist's Tree," undergoes the reaction  $\text{Cu} + 2 \text{Ag}^+ \rightarrow 2 \text{Ag} + \text{Cu}^{2+}$

The copper, Cu, is \_\_\_\_\_ electrons and being \_\_\_\_\_.

- (A) gaining, oxidized  
(B) gaining, reduced  
(C) **losing, oxidized**  
(D) losing, reduced


47. (SKIP) When we reacted  $\text{AgNO}_3$  and  $\text{K}_2\text{CrO}_4$  to form  $\text{Ag}_2\text{CrO}_4$  and  $\text{KNO}_3$ , \_\_\_\_\_ was oxidized.

- (A) chromate ion  
(B) nitrate ion  
(C) silver ion  
(D) **nothing (it wasn't a redox reaction)**

48. What mass of sulfur dioxide,  $\text{SO}_2$  (64.0 g/mol), is produced when 245 g of sulfuric acid,  $\text{H}_2\text{SO}_4$  (98.0 g/mol) reacts completely with zinc metal according to the balanced equation below?  $\text{Zn} + \text{H}_2\text{SO}_4 \rightarrow \text{ZnSO}_4 + \text{H}_2$


- (A) 64.0 g  
(B) **80.0 g**  
(C) 128 g  
(D) 160 g

49. At STP, how many liters of oxygen gas react with 4.00 moles of  $\text{PH}_3$  according to this equation?




- (A) 32.0  
(B) 89.6  
(C) **134**  
(D) 146

50. How many moles of  $\text{FeS}_2$  are required to produce 64 grams of  $\text{SO}_2$  according to the balanced equation below?



- (A) 0.40  
**(B) 0.50**  
 (C) 3.2  
 (D) 4.5


51. Is the reaction below endothermic or exothermic?



$\text{H}_f^\circ$ : +51                  -46      0

- (A) Endothermic  
**(B) Exothermic**  
 (C) Neither endothermic nor exothermic  
 (D) It cannot be determined

52. What is the value of  $\Delta H_{\text{rxn}}$  for the following reaction?



$\text{H}_f^\circ$ : -635    -394    -1207

- (A) -2236 kJ/mol  
**(B) -178 kJ/mol**  
 (C) +178 kJ/mol  
 (D) +2236 kJ/mol

53. **(SKIP)** Consider the following reaction:




The average bond energies in (kJ/mol) are:

$\text{C}\equiv\text{O}$  1070     $\text{C}=\text{O}$  745     $\text{O}=\text{O}$  495

What is the value of  $\Delta H_{\text{rxn}}$ ?

- (A) -1145 kJ/mol  
**(B) -345 kJ/mol**  
 (C) +345 kJ/mol  
 (D) +1145 kJ/mol

54. **(SKIP)** Consider the reaction:



The average N-N bond energy is 160 kJ/mol.

What is the value of  $\Delta H_{\text{rxn}}$ ?

- (A) -160 kJ/mol**  
 (B) 0 kJ/mol  
 (C) +160 kJ/mol  
 (D) +320 kJ/mol


55. Which one of the following changes will result in a decreased rate of reaction?

- (A) adding a catalyst  
 (B) heating up the reactants  
 (C) cutting the reactants into smaller pieces  
**(D) diluting the reactants**

56. Photographic film is sometimes kept in the refrigerator


- (A) because cold film results in sharper pictures.  
**(B) to slow down the chemical reactions on the film.**  
 (C) to protect the film from light.  
 (D) because the cold insulates the film.

57. What distance corresponds to the activation energy for the reaction of X to Y?




- (A) a  
**(B) b**  
 (C) c  
 (D) d

58. What is the  $\Delta H_{\text{rxn}}$  for the reaction shown on the graph?



- (A) -15 kcal/mol  
**(B) -5 kcal/mol**  
 (C) +5 kcal/mol  
 (D) +10 kcal/mol

Questions 59-60: Refer to the following kinetic energy graph. Lines A and B represent the average kinetic energy and the threshold energy, respectively.



59. If the temperature of the system is increased, how will the graph change?

- (A) A will increase, while B remains the same.
- (B) B will increase, while A remains the same.
- (C) Both A and B will increase.
- (D) Both A and B will remain the same.

60. If B decreases while A increases, which of the following may have happened?


- I. The reaction mixture was heated.
- II. A catalyst was added.
- III. The surface area of the reaction mixture was increased.

- (A) I only.
- (B) II and III only.
- (C) I and II only.
- (D) I, II, and III.

61. Consider the reaction:  $2 \text{NO(g)} + \text{O}_2\text{(g)} \rightleftharpoons 2 \text{NO}_2\text{(g)}$   
At equilibrium,  $[\text{NO}] = 0.10 \text{ M}$ ,  $[\text{O}_2] = 0.10 \text{ M}$ , and  $[\text{NO}_2] = 0.010 \text{ M}$ . This reaction is considered:


- (A) Reactant-favored
- (B) Product-favored
- (C) Neither reactant- nor product-favored
- (D) It cannot be determined

62. What is the expression for the equilibrium constant for the following reaction:



- (A)  $\frac{[\text{H}_2][\text{I}_2]}{[\text{HI}]^2}$
- (B)  $\frac{[\text{HI}]^2}{[\text{H}_2][\text{I}_2]}$
- (C)  $\frac{[\text{H}_2]}{[\text{HI}]^2}$
- (D)  $\frac{[\text{HI}]^2}{[\text{H}_2]}$

63. This reaction takes place in solution:




At equilibrium,  $[\text{A}] = 0.1 \text{ M}$  and  $[\text{B}] = 0.1 \text{ M}$ .

What is the value of the equilibrium constant,  $K_{\text{eq}}$ ?

- (A) 0.10
- (B) 0.20
- (C) 1.0
- (D) 10

Questions 64-68: The value of  $K_{\text{eq}}$  for the following reaction is  $1.5 \times 10^{-9}$ .



64. Which of the following may indicate that equilibrium has been reached?

- I.  $[\text{H}_2\text{SO}_3] = [\text{H}^+] = [\text{SO}_3^{2-}]$
- II. Equal moles of  $\text{H}^+$  and  $\text{SO}_3^{2-}$  have been added.
- III. The pH remains constant.

- (A) II only.
- (B) III only.
- (C) I and III only.
- (D) I, II, and III.

65. Which of the following is the correct expression for the equilibrium constant,  $K_{\text{eq}}$ , for the reaction?

- (A)  $\frac{[\text{H}_2\text{SO}_3]}{[\text{H}^+][\text{SO}_3^{2-}]}$
- (B)  $\frac{[\text{H}_2\text{SO}_3]}{[\text{H}^+]^2[\text{SO}_3^{2-}]}$
- (C)  $\frac{[\text{H}^+]^2[\text{SO}_3^{2-}]}{[\text{H}_2\text{SO}_3]}$
- (D)  $\frac{2[\text{H}^+][\text{SO}_3^{2-}]}{[\text{H}_2\text{SO}_3]}$

66. In an equilibrium mixture,  $[\text{H}_2\text{SO}_3] = 0.0015 \text{ M}$ , and  $[\text{SO}_3^{2-}] = 1.0 \times 10^{-2} \text{ M}$ . What is the  $[\text{H}^+]$  in this mixture?

- (A)  $1.5 \times 10^{-5} \text{ M}$
- (B)  $1.0 \times 10^{-4} \text{ M}$
- (C)  $1.0 \times 10^{-2} \text{ M}$
- (D)  $2.0 \times 10^{-2} \text{ M}$

67. To a mixture already at equilibrium, which of the following would have a greater concentration if  $\text{H}_2\text{SO}_3$  (aq) were added?

I.  $\text{H}_2\text{SO}_3$

II.  $\text{H}^+$

III.  $\text{SO}_3^{2-}$

(A) I only.

(B) II only.

(C) II and III only.

**(D) I, II, and III**

68. Which of the following conditions would produce the highest  $[\text{H}_2\text{SO}_3]$ ?


(A) **pH = 2.0**      **T = 10°C**

(B) pH = 2.0      T = 40°C

(C) pH = 9.0      T = 10°C

(D) pH = 9.0      T = 40°C

69. Consider the reaction:



which of the following would shift the reaction to produce more  $\text{SO}_2(\text{g})$ ?


(A) Add  $\text{O}_2$  to the reaction mixture.

**(B) Increase the volume of the container.**

(C) Lowering the temperature.

(D) Remove  $\text{SO}_3(\text{g})$  to the reaction mixture.

70. In the reaction



Under which conditions would produce the most  $\text{Br}_2(\ell)$ ?

(A) High temperature, low pressure

**(B) High temperature, high pressure**

(C) Low temperature, low pressure

(D) Low temperature, high pressure

### Answers

1. C	11. B	21. C	31. D	41. B	51. B	61. A
2. D	12. B	22. C	32. B	42. C	52. B	62. D
3. D	13. D	23. A	33. A	43. C	53. B	63. A
4. D	14. B	24. A	34. A	44. D	54. A	64. B
5. A	15. D	25. A	35. A	45. B	55. D	65. C
6. B	16. D	26. D	36. A	46. C	56. B	66. A
7. B	17. B	27. A	37. C	47. D	57. B	67. D
8. B	18. A	28. D	38. D	48. B	58. B	68. A
9. A	19. C	29. B	39. D	49. C	59. A	69. B
10. D	20. D	30. B	40. C	50. B	60. C	70. B



**Part 2: Free Response.** For each question, write your response in the space provided. If the problem requires mathematical computation, show your work (steps) neatly, reporting your answer with the correct number of significant digits and units. Partial credit is given only when the process taken is clearly shown. Place a box around or circle your final answer.

- Unit conversions should be shown using dimensional analysis, showing how all units cancel out.
- Work for problems involving formulas should follow the I.E.S.A. form.

1. A particular compound containing only chlorine and oxygen is 52.56% chlorine by mass. The molar mass of the compound is found to be between 60 and 70 g/mol.

- a. Find the empirical formula of the compound. (3 points)

$$n_{\text{Cl}} = 52.56 \text{ g} \left( \frac{1 \text{ mol Cl}}{35.45 \text{ g Cl}} \right) = 1.48 \text{ mol Cl}$$

$$n_{\text{O}} = 47.44 \text{ g} \left( \frac{1 \text{ mol O}}{16.00 \text{ g O}} \right) = 2.97 \text{ mol O}$$

$$\text{Cl}_{\frac{1.48}{1.48}} \text{O}_{\frac{2.97}{1.48}} \rightarrow \text{ClO}_2$$

- b. Find the molecular formula of the compound. (2 points)

$$\text{Mass of Empirical Formula: } 1(35.45) + 2(16.00) = 67.45 \text{ g/mol}$$

$$\frac{\text{Molar Mass}}{\text{Empirical Formula Mass}} = \frac{60-70 \text{ g/mol}}{67.45 \text{ g/mol}} \approx 1$$

$$\text{Molecular Formula} = 1(\text{ClO}_2) = \text{ClO}_2$$

- c. A 1.680 L sample of this compound was obtained at STP. How many molecules are in this sample? (3 points)

$$1.680 \text{ L} \left( \frac{1 \text{ mol}}{22.4 \text{ L}} \right) \left( \frac{6.02 \times 10^{23} \text{ molecules}}{1 \text{ mol}} \right) = 4.52 \times 10^{22} \text{ molecules}$$

2. Consider the compounds CO and CaO. Draw the Lewis Structures for the following compounds. Then identify the type of solid formed, and the major inter-particle forces involved. (3 points)


Compound:	CO	CaO
Lewis Structures:	$\text{:C} \equiv \text{O:}$	$\left[ \text{Ca} \right]^{2+} \left[ \text{:}\ddot{\text{O}}\text{:} \right]^{2-}$
Type of Solid Formed:	<b>Molecular Solid</b>	<b>Ionic Solid</b>
Inter-Particle Forces	<b>Dipole-dipole Interactions</b>	<b>Ionic Bonds</b>

3. Complete the chart for the following compounds: (5 points)


Compound:	BI <sub>3</sub>	SF <sub>2</sub>
Lewis Structure	<pre>       :I:         :I: B :I:               :I:           </pre>	<pre>       :S: :F:         :       :F:           </pre>
Molecular Shape	<b>Trigonal Planar</b>	<b>Bent</b>
Bond Angles	<b>120°</b>	<b>104°</b>
Molecule is	<b>Polar   Non-Polar</b>	<b>Polar   Non-Polar</b>
Type of IMF:	<b>London Dispersion Forces</b>	<b>Dipole-dipole Interactions</b>

4. A refrigerant has a molecular formula C<sub>4</sub>H<sub>10</sub>.


a. Draw two possible structural formulas for C<sub>4</sub>H<sub>10</sub> and name them (**SKIP NAMING**). (4 points)



b. Write the balanced equation for the combustion of C<sub>4</sub>H<sub>10</sub>. (4 points)



5. Write the equation for the reaction between propene and chlorine gas using structural formulas. (2 points)



6. Complete the table for aqueous samples of each compound, predicting its color in cabbage juice and its conductivity with a light bulb, and writing its dissociation equation. (4 points)

Compound	Cabbage Juice Color [red   green   purple]	Conductivity [bright   dim   dark]	Dissociation Equation
(i) HBr	<b>RED</b>	<b>BRIGHT</b>	$\text{HBr} \rightarrow \text{H}^+ + \text{Br}^-$
(ii) Mg(OH) <sub>2</sub>	<b>GREEN</b>	<b>DIM</b>	$\text{Mg(OH)}_2 \rightleftharpoons \text{Mg}^{2+} + 2 \text{OH}^-$
(iii) NH <sub>3</sub>	<b>GREEN</b>	<b>DIM</b>	$\text{NH}_3 + \text{H}_2\text{O} \rightleftharpoons \text{NH}_4^+ + \text{OH}^-$

7. Find the pH of a 0.350 M HC<sub>3</sub>H<sub>6</sub>O<sub>2</sub> solution. K<sub>a</sub> of HC<sub>3</sub>H<sub>6</sub>O<sub>2</sub> is  $1.3 \times 10^{-5}$ . Show your work using a dissociation equation, K<sub>a</sub> expression, and an ICE Table. (4 points)


HC <sub>3</sub> H <sub>6</sub> O <sub>2</sub> ⇌ H <sup>+</sup> + C <sub>3</sub> H <sub>6</sub> O <sub>2</sub> <sup>-</sup>				$K_a = \frac{[H^+][C_3H_6O_2^-]}{[HC_3H_6O_2]} = 1.3 \times 10^{-5}$ $\frac{(x)(x)}{(0.35 - x)} = 1.3 \times 10^{-5}$ $x = 2.1 \times 10^{-3} = [H^+]$ $pH = -\log(2.1 \times 10^{-3}) = 2.67$
Initial	0.350 M	0 M	0 M	
Change	- x	+ x	+ x	
Equil.	0.35 - x	x	x	

8. Consider the following reactions. Lead metal is placed in a solution of hydrochloric acid. (3 points)

- a. Balanced molecular equation: **Pb (s) + 2 HCl (aq) → H<sub>2</sub> (g) + PbCl<sub>2</sub> (s)**
- b. Balanced net ionic equation. **Pb (s) + 2 H<sup>+</sup> + 2 Cl<sup>-</sup> → H<sub>2</sub> (g) + PbCl<sub>2</sub> (s)**
- c. **(SKIP)** Which element is **Pb** Which element is reduced? **H (or H<sup>+</sup>)**  
oxidized?

9. Consider the compound isopropyl alcohol, or 2-propanol, C<sub>3</sub>H<sub>8</sub>O (ℓ).

- a. Write the balanced equation for the combustion of 2-propanol. (1 point)



- b. What volume (in L) of CO<sub>2</sub> at STP can be produced if 4.05 g C<sub>3</sub>H<sub>8</sub>O is burned? (3 points)

$$\text{Molar Mass} = 3(12.01) + 8(1.008) + 1(16.00) = 60.094 \text{ g/mol}$$

$$4.05 \text{ g C}_3\text{H}_8\text{O} \left( \frac{1 \text{ mol C}_3\text{H}_8\text{O}}{60.094 \text{ g C}_3\text{H}_8\text{O}} \right) \left( \frac{6 \text{ mol CO}_2}{2 \text{ mol C}_3\text{H}_8\text{O}} \right) \left( \frac{22.4 \text{ L CO}_2}{1 \text{ mol CO}_2} \right) = 4.53 \text{ L CO}_2$$

- c. If 4.25 L of CO<sub>2</sub> was actually produced in (b) above, what was the percent yield of CO<sub>2</sub>? (1 point)

$$\% \text{ yield} = \frac{\text{actual yield}}{\text{theoretical yield}} \times 100\% = \frac{4.25 \text{ L}}{4.53 \text{ L}} \times 100\% = 93.8\%$$

- d. What is the ΔH<sub>rxn</sub> if 1025 kJ of energy is released when 30.5 g C<sub>3</sub>H<sub>8</sub>O is burned? (2 points)

$$\Delta H_{\text{rxn}} = \frac{q}{n_{\text{rxn}}} = \frac{-1025 \text{ kJ}}{30.5 \text{ g C}_3\text{H}_8\text{O}} \left( \frac{60.094 \text{ g C}_3\text{H}_8\text{O}}{1 \text{ mol C}_3\text{H}_8\text{O}} \right) \left( \frac{2 \text{ mol C}_3\text{H}_8\text{O}}{1 \text{ mol reaction}} \right) = -4039 \text{ kJ/mol}$$

- e. **(SKIP)** The enthalpies of formation, ΔH<sub>f</sub>, of CO<sub>2</sub> (g) and H<sub>2</sub>O (ℓ) are -394 kJ/mol and -286 kJ/mol, respectively. Find the ΔH<sub>f</sub> of C<sub>3</sub>H<sub>8</sub>O. (2 points)

$$\Delta H_{\text{rxn}} = [\Delta H_{\text{f,products}}] - [\Delta H_{\text{f,reactants}}]$$

$$-4039 \text{ kJ/mol} = [(6 \text{ mol CO}_2)(-394 \text{ kJ/mol CO}_2) + (8 \text{ mol H}_2\text{O})(-286 \text{ kJ/mol H}_2\text{O})] -$$

$$[(2 \text{ mol C}_3\text{H}_8\text{O})(\Delta H_{\text{f,C}_3\text{H}_8\text{O}}) + (9 \text{ mol O}_2)(0 \text{ kJ/mol O}_2)]$$

$$\Delta H_{\text{f,C}_3\text{H}_8\text{O}} = 306.5 \text{ kJ/mol}$$



The following reaction rate data was obtained:

Trial	$[\text{NH}_3]$	$[\text{H}_2\text{O}_2]$	Initial Rate of Appearance of $\text{N}_2\text{H}_4$
1	0.010 M	0.010 M	3.6 M/min
2	0.010 M	0.030 M	10.8 M/min
3	0.020 M	0.010 M	14.4 M/min

- a. Write the rate law for this reaction. (3 points)

$$\text{Rate} = k[\text{NH}_3]^x[\text{H}_2\text{O}_2]^y$$

Using experiments 1 and 3,

$$\frac{14.4 \text{ M/min}}{3.6 \text{ M/min}} = \frac{k(0.020 \text{ M})^x(0.010 \text{ M})^y}{k(0.010 \text{ M})^x(0.010 \text{ M})^y} \quad 4 = (2)^x \quad x = 2$$

Using experiments 1 and 2,


$$\frac{10.8 \text{ M/min}}{3.6 \text{ M/min}} = \frac{k(0.010 \text{ M})^2(0.030 \text{ M})^y}{k(0.010 \text{ M})^2(0.010 \text{ M})^y} \quad 3 = (3)^y \quad y = 1$$

$$\text{Rate} = k[\text{NH}_3]^2[\text{H}_2\text{O}_2]^1$$

- b. Find the value of the rate constant, k. Include units. (1 points)

Using Experiment 1

$$k = \frac{\text{rate}}{[\text{NH}_3]^2[\text{H}_2\text{O}_2]^1} = \frac{3.6 \text{ M/min}}{(0.010 \text{ M})^2(0.010 \text{ M})^1} = 3.6 \times 10^6 \text{ M}^{-2} \text{ min}^{-1}$$



- a. Write the equilibrium constant expression for this reaction. (1 point)

$$K_{\text{eq}} = \frac{[\text{HCl}]_{\text{eq}}^2}{[\text{H}_2]_{\text{eq}}[\text{Cl}_2]_{\text{eq}}}$$

- b. Experiment #1: A reaction mixture is found to contain the following:


$$[\text{H}_2] = 0.020 \text{ M}, [\text{Cl}_2] = 0.020 \text{ M}, [\text{HCl}] = 0.38 \text{ M}$$

In which direction will the reaction shift? Justify with calculations. (3 points)

$$Q = \frac{[\text{HCl}]^2}{[\text{H}_2][\text{Cl}_2]} = \frac{(0.38)^2}{(0.020)(0.020)} = 361 < 841$$

Because  $Q < K_{\text{eq}}$ , the reaction will go in the forward direction.

- c. What are the concentrations of all substances when the reaction reaches equilibrium? Show work using an ICE Chart. (4 points)



$$\text{I} \quad 0.020 \quad 0.020 \quad 0.38$$

$$\text{C} \quad -x \quad -x \quad +2x$$

$$\text{E} \quad 0.02-x \quad 0.02-x \quad 0.38+2x$$

$$K_{\text{eq}} = \frac{[\text{HCl}]_{\text{eq}}^2}{[\text{H}_2]_{\text{eq}}[\text{Cl}_2]_{\text{eq}}} = \frac{(0.38+2x)^2}{(0.020-x)(0.020-x)} = 841 \quad x = 0.00645 \text{ M}$$

$$[\text{H}_2]_{\text{eq}} = [\text{Cl}_2]_{\text{eq}} = 0.014 \text{ M}; [\text{HCl}]_{\text{eq}} = 0.393 \text{ M}$$